

# **FOR CRABGRASS CONTROL (COASTAL SOUTH) - HIGH CUT TURFGRASS (PROGRAM 1)**

## General Information

### PRODUCT INFORMATION

This product contains turf fertilizer plus a selective herbicide for control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turf. Apply this product only to seeded, sodded, or sprigged turf that is well established. Turf must develop a good root system and uniform stand, and have received at least two mowings following seeding, sodding or sprigging before making the first application of this product. Use of this product on turf that is not well established, or weakened by weather, pests or other stress, may increase the chance of turf injury.

This product provides pre-emergence control of crabgrass (large, smooth, and southern species) when applied prior to emergence of crabgrass from the ground. In addition, this product provides “early post-emergence” control during the early stages of growth after crabgrass emerges from the ground. This early post-emergence crabgrass control is limited, and post-emergence applications are only effective if applied before crabgrass has displayed its fifth leaf or first tiler (which generally corresponds to time when you can first see the crabgrass plants in the lawn). The benefit of this post-emergence activity is that it gives the user a 2 to 4 week longer period of time to make applications for effective control of crabgrass.

### USE PRECAUTIONS

Do not apply more than 8.8 pounds per 1000 square feet (0.5 pounds AI per acre) per application. Do not apply more than 26.4 pounds per 1000 square feet (1.5 pounds AI per acre) per year. (In New York, do not apply more than 8.8 lbs per 1000 square feet per year.)

Cultural practices such as aerification and de-thatching should be completed, and the turf allowed to recover before application of this product. Do not use clippings from treated turf for animal feed or for mulching around vegetables or fruit trees. Do not allow domestic animals to feed or forage on treated turf. Keep people and

pets off treated areas until dust has settled. Do not use on golf course putting greens.

#### TIPS FOR IMPROVED CONTROL

For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. Irrigation or rainfall within 7 days after application may improve weed control. Use of split (half rate) applications spaced 6-8 weeks apart may provide improved weed control.

#### RESEEDING, OVERSEEDING, OR SPRIGGING:

Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When seeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, use "slit seeders" for good seed-to-soil contact.

Refer to the product label for the Spreader setting to apply at the listed rate (lbs per 1000 sq ft).

#### Limitations, Restrictions, and Exceptions

#### FOR CRABGRASS CONTROL (COASTAL SOUTH) - HIGH CUT TURFGRASS (PROGRAM 1)

##### Crops:

Creeping Bentgrass - Turfgrass injury may result from use of this product on certain varieties of creeping bentgrass, such as Cohansey, Carmen, Seaside and Washington. Not all varieties of creeping bentgrass have been tested. Do not apply this product to Colonial Bentgrass (*Agrostis tenuis*) varieties.

Fine Fescue - Turfgrass injury may result from use of this product on the following Fescue varieties which are sensitive to this product: Atlanta, Banner, Beauty, Bilgart, CF-2, Enjoy, HF-93, Highlight, Ivalo, Koket, Jamestown, Majenta, Mary, Pennlawn, Tamara, Tatjana, Walfort and Waldina. Not all varieties of fescue have been tested.

Buffalograss - Do not use this product on seedling Buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

SPRING APPLICATIONS: For single applications in the spring or early summer, use the appropriate rate from “Application Rates” table (depending on user’s location, mowing height, pre- or post-emergent use). The duration of residual weed control is directly related to the total rate applied, but varies depending on weather, weed pressure and location. Use “Program 1” for preemergence control on areas where turf is cut high (home lawns) for 3 to 5 months of crabgrass control. This program should not be used for postemergence crabgrass control. Use “Program 2” for preemergence control where turf is cut low (golf course fairways) or where weed control was not conducted the previous year. This program provides 4 to 6 months of preemergence weed control, and also for early postemergence control of crabgrass. This product may be applied as a single application or as a split application for crabgrass control in the spring, summer or fall.

Method

[N. A.](#)

Rates

[field rates 0](#)

[field rates 1](#)

[field rates 2](#)

•

Timings

[Preemergence \(Weed\)](#)

[In the spring or early summer.](#)