

# **FOR CONTROL OF HOUSE MOUSE**

## General Information

### USE RESTRICTIONS

This product may only be used to control Norway rats, roof rats and house mice in and around agricultural buildings. Do not use in homes or other human residences. Do not apply further than 50 feet from agricultural buildings. Do not sell this product in individual containers holding less than 8 pounds of bait. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment. Do not broadcast bait.

### Selection of Treatment Areas:

Determine areas where rats and/or house mice will most likely find and consume bait. Generally, these are along walls, by gnawed openings, in corners and concealed places, between floors or walls, in or beside burrows, or in locations where rats and/or house mice or their signs have been observed. Remove as much alternative food as possible.

### APPLICATION DIRECTIONS:

Follow-up: Replace contaminated or spoiled bait immediately. Wearing waterproof gloves, collect and dispose of all visible dead animals and leftover bait, as well as any bait that has become accessible to non-target animals. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. For a continuous infestation, set up permanent bait stations and replenish bait as needed.

### Limitations, Restrictions, and Exceptions

### FOR CONTROL OF HOUSE MOUSE

Apply 1/4 to 1/2 ounce per placement, usually spaced 8 to 12 feet apart. Up to 2 ounces may be needed at points of very high house mouse activity. Maintain a constant supply of fresh bait for 15 days or until signs of house mouse activity cease.

Method

[Bait](#)

Rates

[field\\_rates 0](#)

- 

Timings

[Where rats and/or house mice or their signs have been observed.](#)