

TURFGRASS - SUPPRESSION OF FUSARIUM BLIGHT, NECROTIC RING SPOT, SUMMER PATCH

Limitations, Restrictions, and Exceptions

TURFGRASS

How to Use

- Apply with a drop-type or rotary spreader. Do not apply by hand or hand-held rotary spreader. For best results with a rotary spreader, apply during a calm period of the day. If necessary to apply when a breeze is blowing, move in the direction of the wind to prevent material from blowing toward the operator.
- Allow one week between application of this and other control products. (This guideline may be relaxed where a severe disease attack requires immediate treatment.)
- Maximum number of applications is four per year.
- It is not necessary to water-in immediately. However, water is required before product will be activated.

When to Use

Prevention

- Apply to dry or moist foliage at the PREVENTIVE RATE when disease activity is noted or suspected.
- Repeat at 2- to 3-week intervals. During periods of high day and night temperature and humidity, two week treatment interval may be needed to achieve best brown patch control.
- For cure of these diseases when symptoms indicate advanced stages, apply at the NORMAL RATE (CURATIVE) and follow with the PREVENTIVE RATE within 2 -3 weeks.

Suppressing turfgrass diseases

- Apply to dry foliage at the NORMAL RATE (CURATIVE) every 30 days (beginning in

late spring to early summer).

- Water-in immediately after application and keep turf from becoming drought stressed by watering (1 inch) weekly.
- Three to four applications may be necessary for season-long suppression.

Do not apply more than 2.7 pounds per 1,000 sq ft. of turf.

Method

Broadcast

Timings

Every 30 days (beginning in late spring to early summer).