

FOR CONTROL OF LARGER ANNUALS AND BURNDOWN OF PERENNIAL WEED GROWTH

General Information

Weed Pharm is a fast-acting, non-selective contact herbicide containing an acid for which the residues degrade rapidly in the soil. Foliar contact results in rapid desiccation and control of annual weeds and grasses, and top growth reduction of herbaceous perennial weeds and grasses (see tables below). Retreatment may be required for control of established perennial weeds. Weed Pharm can be applied up to and including the day of harvest.

Weed Pharm Weed & Grass Killer is a 20.0% Acetic Acid herbicide labeled for use as a horticultural spray for residential use, farmstead, food crops, non-production agricultural use, institutional use, and users at all sites indicated below.

DO NOT SPRAY DIRECTLY ONTO FOOD CROPS

Use Weed Pharm to control weeds which surround all food crops, non-food crops (ornamentals) and non-production agricultural, farmstead, right-of-way, and institutional land sites. Use to control weeds in interstate freeways/highways, federal and state highways and city/county roads and walk ways, railroad rights-of-way, tank farms, power stations and easements, and other rights-of-way, industrial sites, and vacant lots. Suitable to keep down weeds on all botanical and private association ornamental gardens, kennels, dog runs, and other animal enclosures and walk ways.

Limitations, Restrictions, and Exceptions

FOR CONTROL OF LARGER ANNUALS AND BURNDOWN OF PERENNIAL WEED GROWTH

Larger annual weeds and perennials are more difficult to control and may require retreatment. Treat initially as recommended above and repeat if new growth of leaves appears. Thorough coverage of all weed foliage is necessary to achieve desirable control. To reduce amount of product required controlling well-established weed stands, mowing to height of 4-6 inches and remove cut vegetation before

application of product.

TIMING OF APPLICATIONS:

Best results are achieved from applications to actively growing young weeds. Weeds that are mature, dormant or hardened due to moisture stress are more tolerant of herbicide treatments. For best results spray in full sunshine at temperatures above 65°F. Only contracted vegetation will be affected.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

Liverworts - For liverworts dilute Weed Pharm at a rate of 1 gallon of Weed Pharm to 2 gallons of water. For all other weeds use full strength.

Method

[Spray](#)

Rates

[field_rates 0](#)

-

Timings

[Best results are achieved from applications to actively growing young weeds.](#)