

FOOD AND FEED-HANDLING AREAS - BROADCAST (LOW)

General Information

PRODUCT INFORMATION

Zyrox Fly Granular Bait contains the active ingredient cyantraniliprole in a granular bait formulation for use to control house flies. A fly control program includes identification, inspection, sanitation, mechanical control and control using an insecticide product depending on the conditions and use site. Fly control baits such as Zyrox Fly Granular Bait can be an important tool in the management of fly populations as part of an integrated pest management program.

- Ready-to-use granular bait contains 0.5% cyantraniliprole.
- Controls house flies in and around the outside of structures and protected or enclosed areas.
- Alternative house fly control chemistry for use in IPM and resistance management programs.

APPLICATION USE SITES

- Outdoor areas used for storage of waste in refuse dumpsters that are fenced in or inaccessible to children and pets around commercial establishments such as restaurants, taverns, hotels, grocery stores, supermarkets, bakeries, warehouses, and sports complexes.
- Outdoor areas of commercial operations such as, but not limited to canneries, beverage processing plants, dairy, meat and poultry processing plants, fruit and vegetable processing plants, seafood processing plants, tanneries, rendering plants, grease collection stations, kennels, schools, recycling plants, and military installations.
- Outdoor areas of agricultural animal production facilities such as poultry/broiler houses, walkways of caged layer houses, swine production structures, livestock housing structures, and horse stables.
- Use in indoor areas at all the above described use sites is permitted only if is

placed in bait stations, with the exception of walkways within caged layer houses. Refer to Use in Bait Stations section for specific use instructions.

Food/Feed Handling Establishments

(Food/feed handling establishments are places other than private residences in which food/feed is held, processed, prepared or served.)

Specific Restrictions

- Do not make broadcast applications of Zyrox Fly Granular Bait in indoor areas of residential, institutional, public, agricultural buildings/structures and food/feed handling establishments.
- For indoor areas of residential, institutional, public, agricultural buildings/structures, or food/feed handling establishments, Zyrox Fly Granular Bait must be placed in commercially available bait stations designed for granular fly control baits. Indoor use as a scatter bait is permitted only in caged layer poultry houses.
- Do not apply to areas or surfaces where food/feed, food utensils or food processing surfaces may come into contact and become contaminated.

USE RESTRICTIONS

- Bait stations must be used when using Zyrox Fly Granular Bait inside single or multi-unit residential structures, commercial establishments, food processing plants, restaurants and animal production facilities.
- When bait stations are being used, place them where the stations are inaccessible to children or pets.
- Do not place bait in areas accessible to food producing animals or animals being grown for food.
- Do not place bait in areas accessible to children and pets.
- Do not use in food/feed areas of food/feed handling establishments unless placed in bait stations designed to hold granular baits.
- Not for use in edible product areas of food processing plants or other areas where

food is commercially prepared or processed unless placed in bait stations designed to hold granular baits.

- For outdoor broadcast applications, do not apply more than 2 lb bait/1,000 sq ft/year (equivalent to 0.01 lb ai/1,000 sq ft/year or 0.44 lb ai/Acre/year).

- Do not apply in outdoor areas when raining.

Limitations, Restrictions, and Exceptions

Applications in Food/Feed Handling Areas

Application to food/feed areas of food/feed handling establishments may only be made using commercially available bait stations designed for use of granular fly control baits.

Food and feed handling areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered a food/feed area when food is exposed and facility is in operation.

APPLICATION PROCEDURES AND USE RATES

Broadcast

Broadcast or scatter Zyrox Fly Granular Bait using suitable granular spreader equipment or directly from the container evenly to the target areas. Avoid placing the fly bait granules in piles and scatter bait as evenly as possible over surfaces.

Reapply bait every 7 days until fly population is at acceptable levels. For outdoor broadcast applications, DO NOT apply more than 2 lb/1,000 sq ft per calendar year.

Method

[Broadcast](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Timings

[When fly populations are low.](#)