

# HOUSE MICE

## General Information

### DIRECTIONS FOR USE

**IMPORTANT:** Do not expose children, pets, or other non-target animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled and unconsumed bait as specified on the label.

**USE RESTRICTIONS:** For control of Norway rats, roof rats and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures. This product is especially suited for use in wet or damp areas, including river banks, gullies, irrigation ditches, sewers, garbage dumps and landfills.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not apply bait directly to ground surface or in grass or other ground cover.

Wear protective gloves when applying bait. With detergent and hot water, wash all implements used for applying bait. Do not use these implements for mixing, holding or transferring foods or feeds.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed

openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

#### FOLLOW-UP:

- Replace contaminated or spoiled bait immediately. Using waterproof gloves collect and dispose of all dead, exposed animals and leftover bait properly. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

#### Limitations, Restrictions, and Exceptions

#### HOUSE MICE:

Break Ramik Bar into discrete 1-ounce. pieces. Apply one 1-ounce piece per placement. Space placements at 8- to 12-foot intervals. Placement of two 1-ounce pieces each may be needed at points of very high house mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until there no longer are signs of new feeding by house mice.

#### Method

[Bait](#)

Timings

[N.A.](#)