

## **FRUITS AND NUTS: ALMONDS**

General Information

### **DIRECTIONS FOR USE**

For household use only.

**SHAKE WELL BEFORE USING.**

**KEEP FROM FREEZING.**

The control of fungal disease is based on prevention, including keeping garden and border areas clean, destroying leaves in the fall, and applying fungicides either before or at the onset of disease. Plant surfaces must be completely covered with fungicide to successfully prevent infection. Repeat applications may be needed for adequate control. Use the highest rate indicated during conditions of severe disease pressure, such as wet or humid weather. The lower rate is suitable for general preventive applications.

Limitations, Restrictions, and Exceptions

### **FRUITS AND NUTS**

Mix specified amount of KOP-R-SPRAY in one gallon water. Apply at the rate of one gallon mixed spray to 200 sq. ft. of garden area or up to 2 gallons mixed spray to a large tree. Thoroughly wet foliage to the point of runoff, taking care to cover all plant surfaces. Agitate the container of mixed spray regularly during application.

**ALMONDS:** Use 3 to 4 tablespoons to control Shothole. Apply at leaf fall to protect buds and shoots from infection during rainy periods. Reapply every 3 to 4 weeks up until late bud swell — the period in spring before buds begin to open. Do not apply after full bloom. Use 3 to 4 tablespoons to control Brown Rot. Apply at delayed dormant bud swell stage.

Delayed Dormant — The period just before blossom buds break open.

Method

[Spray](#)

Rates

[field\\_rates 0](#)

- 

Timings

[Brown Rot: At delayed dormant bud swell stage.](#)

[Shothole: At leaf fall to protect buds and shoots from infection during rainy periods.](#)