

## **VEGETABLES: BEANS (SNAP AND DRY)**

General Information

### DIRECTIONS FOR USE

For household use only.

### SHAKE WELL BEFORE USING KEEP FROM FREEZING

The control of fungal disease is based on prevention, including keeping garden and border areas clean, destroying leaves in the fall, and applying fungicides either before or at the onset of disease. Plant surfaces must be completely covered with fungicide to successfully prevent infection. Repeat applications may be needed for adequate control. Use the highest rate indicated during conditions of severe disease pressure, such as wet or humid weather. The lower rate is suitable for general preventive applications.

Limitations, Restrictions, and Exceptions

### VEGETABLES

Mix specified amount of KOP-R-SPRAY in one gallon water. Apply at the rate of one gallon mixed spray to 200 sq. ft. of garden area. Thoroughly wet foliage to the point of runoff, taking care to cover all plant surfaces. Shake sprayer often during application.

BEANS (SNAP AND DRY): Use 1 to 3 teaspoons to control Bacterial Blight and Halo Blight. Begin when plants are 3 to 5 inches high and before diseases appear. Repeat at 7 to 10 day intervals or at 5 to 7 day intervals under conditions of severe disease pressure.

Method

[Spray](#)

Rates

[field\\_rates 0](#)

•

Timings

When plants are 3 to 5 inches high and before diseases appear.