

VEGETABLES: BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER - BLACK ROT AND BLACK LEAF SPOT

General Information

DIRECTIONS FOR USE

For household use only.

SHAKE WELL BEFORE USING KEEP FROM FREEZING

The control of fungal disease is based on prevention, including keeping garden and border areas clean, destroying leaves in the fall, and applying fungicides either before or at the onset of disease. Plant surfaces must be completely covered with fungicide to successfully prevent infection. Repeat applications may be needed for adequate control. Use the highest rate indicated during conditions of severe disease pressure, such as wet or humid weather. The lower rate is suitable for general preventive applications.

Limitations, Restrictions, and Exceptions

VEGETABLES

Mix specified amount of KOP-R-SPRAY in one gallon water. Apply at the rate of one gallon mixed spray to 200 sq. ft. of garden area. Thoroughly wet foliage to the point of runoff, taking care to cover all plant surfaces. Shake sprayer often during application.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: Use 2 teaspoons to control Black Rot (*Xanthomonas*) and Blackleaf Spot (*Alternaria*). Apply at 7 to 10 day intervals. Caution: A slight reddening of older leaves may occur on broccoli and a slight flecking of outer leaves may occur on cabbage at this rate.

For control of diseases of these crops, begin applications after transplants are set in the garden, or shortly after the emergence of seedlings, or when conditions favor disease development.

Method

[Spray](#)

Rates

[field_rates 0](#)

-

Timings

[At 7 to 10 day intervals.](#)