

VEGETABLES: TOMATOES

General Information

DIRECTIONS FOR USE

For household use only.

SHAKE WELL BEFORE USING KEEP FROM FREEZING

The control of fungal disease is based on prevention, including keeping garden and border areas clean, destroying leaves in the fall, and applying fungicides either before or at the onset of disease. Plant surfaces must be completely covered with fungicide to successfully prevent infection. Repeat applications may be needed for adequate control. Use the highest rate indicated during conditions of severe disease pressure, such as wet or humid weather. The lower rate is suitable for general preventive applications.

Limitations, Restrictions, and Exceptions

VEGETABLES

Mix specified amount of KOP-R-SPRAY in one gallon water. Apply at the rate of one gallon mixed spray to 200 sq. ft. of garden area. Thoroughly wet foliage to the point of runoff, taking care to cover all plant surfaces. Shake sprayer often during application.

TOMATOES: Use 1 to 3 teaspoons to control Early and Late Blight, Bacterial Speck and Bacterial Spot. Begin applications upon emergence of seedlings or immediately after transplanting. Apply at 5 to 7 day intervals. When disease is severe, apply at 4 to 5 day intervals. Complete coverage is essential for disease control.

Method

[Spray](#)

Rates

[field_rates 0](#)

•

Timings

Begin applications upon emergence of seedlings or immediately after transplanting.