

ORNAMENTALS

General Information

DIRECTIONS FOR USE

For household use only.

SHAKE WELL BEFORE USING KEEP FROM FREEZING

The control of fungal disease is based on prevention, including keeping garden and border areas clean, destroying leaves in the fall, and applying fungicides either before or at the onset of disease. Plant surfaces must be completely covered with fungicide to successfully prevent infection. Repeat applications may be needed for adequate control. Use the highest rate indicated during conditions of severe disease pressure, such as wet or humid weather. The lower rate is suitable for general preventive applications.

Limitations, Restrictions, and Exceptions

ORNAMENTALS

Mix 3 tablespoons KOP-R-SPRAY in one gallon water. Apply at the rate of one gallon mixed spray to 200 sq. ft. of garden area or up to 2 gallons mixed spray to a large tree. Thoroughly wet foliage to the point of runoff, taking care to cover all plant surfaces. Shake the sprayer often during application.

SPECIFIC USE DIRECTIONS FOR ORNAMENTALS

KOP-R-SPRAY is best used as a preventive treatment. Apply before disease appears when disease conditions occur, such as during hot, humid weather or cool, moist weather. Repeat at 7 to 10 day intervals and after rain.

- For anthracnose on sycamore, apply when leaf buds begin to open. Repeat as full coverage spray 10 to 14 days later.
- For bacterial blight of lilac, apply in September and again before fall rains.
- For bacterial leaf spot on barberry, apply when leaves appear. Repeat 2 to 3 times

10 days apart.

- For black canker of willow, apply when new leaves are one-quarter inch long. Repeat in 2 weeks.
- For botrytis blight of peonies, apply before shoots are 1 foot tall. Repeat in 2 weeks, and again if any signs of disease appear.
- For cedar-apple rust of cedar and juniper, apply in July or August.
- For leaf blister of elm and oak, apply before growth begins or as leaves uncurl.
- For leaf gall of azalea, apply to entire plant before buds break in spring. Repeat 2 to 3 weeks later.
- For anthracnose, false smut, leaf spots or scab of palm, apply until stems and leaves are wet at first sign of disease.
- For lichen control, apply as a directed spray to thoroughly wet lichens. Repeat as needed.

Method

[Spray](#)

Rates

[field rates 0](#)

•

Timings

[Before disease appears when disease conditions occur, such as during hot, humid weather or cool, moist weather.](#)