

CILANTRO, CORIANDER - APHIDS, ETC.

General Information

DIRECTIONS FOR USE

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

RESISTANCE MANAGEMENT

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product or other products with a similar mode of action may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

APPLICATIONS INSTRUCTIONS - FOOD CROPS

Application rates will vary according to pest pressure, timing of sprays, and field scouting. Use lower rates under light to moderate infestations and higher rates under heavy insect pressure and for mite control. Arid climates generally require higher rates.

Cultivation within 10 feet of a water body is prohibited to allow for the growth of a vegetated filter strip. In New York State, this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

CHEMIGATION USE DIRECTIONS

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

For LEPA irrigation, a minimum of 0.75 inch of water per acre is recommended. Where non-emulsified oils are used as the diluent, 1 to 2 pints per acre is recommended.

Results from utilizing chemigation have been variable and depend upon the set up and calibration of equipment. Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water. Contact your State Agricultural Extension Service specialists, equipment manufacturers, or other experts for consultation on the suitability of the equipment set up to obtain effective control of the target insect pests.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Failure to cease application during a mechanical stoppage may result in undesirable residues to adjacent area.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain function interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

REVEAL ENDURX should be applied continuously for the duration of the water application. REVEAL ENDURX should be diluted in sufficient volume to ensure accurate application over the area to be treated. When using chemigation, a minimum of 0.5 inch per acre of irrigation water is recommended. Agitation generally is not required when a suitable diluent is used. A diluent test should be conducted to ensure that phase separation will not occur during dilution and application. Failure to achieve a uniform dilution throughout the time of application may result in undesirable residues or less than desirable control.

SPRAY DRIFT REQUIREMENTS

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, ESTUARIES, AND COMMERCIAL FISH FARM PONDS.

Do not apply by ground equipment within 25 feet, or by air within 150 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds. Increase the buffer zone to 450 feet when ultra low volume (ULV) application is made in cotton. Use of ultra low volume (ULV) application on corn and hops is prohibited.

Wind Direction and Speed

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Only apply this product if the wind direction favors on-target deposition. Do not apply when the wind velocity exceeds 15 mph.

Temperature Inversion

Do not make aerial or ground applications into temperature inversions.

Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Do not make aerial or ground applications to corn if heavy rainfall is imminent.

Droplet Size

Use only Medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASAE (S572) definition for standard nozzles. Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.

Ground Applications

For ground applications, wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.

For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

Aerial Applications

The spray boom must be mounted on the aircraft as to minimize drift caused by

wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter.

Flight speed and nozzle orientation must be considered in determining droplet size.

Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

ROTATIONAL CROPS

Crops with existing bifenthrin tolerances may be rotated at any time. All other crops may be rotated 30 days following the final application of REVEAL ENDURX.

Refer in the label for REVEAL ENDURX LAWN DILUTION CHART.

Attention

- Do not apply to pets, crops, or sources of electricity.
- Firewood is not to be treated.
- Do not allow spray to contact food, foodstuffs, food contacting surfaces, food utensils or water supplies.
- Do not apply this pesticide in livestock buildings (barns).
- Keep children and pets off treated areas following application until the spray has dried.
- Do not apply by air.
- Do not use in greenhouses.
- Do not apply this product through any type of irrigation system. Do not apply when a temperature inversion exists.
- Do not apply for surface feeding pests if rain is expected within 12 hours (or whatever time is necessary for the spray to dry).
- For turf treatment, apply with nozzles not more than 2 feet above the grass.
- Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- Do not apply when grass areas are water logged or the soil is saturated with water (i.e., will not accept irrigation).
- Vinyl and Aluminum Siding: Do not spray directly onto vinyl or aluminum siding. If

REVEAL ENDURX inadvertently contacts vinyl and aluminum siding (particularly lightly colored, aged, weathered or otherwise damaged), it may result in staining, bleaching or discoloration. Wash off thoroughly with detergent and water. Factors such as extreme heat and direct sunlight can promote damage when using emulsifiable concentrates. Avoid application to vinyl or aluminum siding while exposed to direct sunlight or during the heat of the day.

Limitations, Restrictions, and Exceptions

CILANTRO, CORIANDER

APPLICATION INSTRUCTIONS

Apply using sufficient water to obtain uniform coverage. Apply as needed. Apply with ground equipment using a minimum of 10 gallons of finished spray per acre or a minimum of 2 gallons per acre by aircraft.

RESTRICTIONS

- Do not make applications less than 7 days apart.
- Do not apply more than 0.5 pound active ingredient per acre per season.
- Do not apply within 3 days of harvest.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Directed](#)

Pre-Harvest Interval

3 days

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

N.A.