

PEANUTS (SOILBORNE DISEASE)

General Information

GENERAL PRECAUTIONS AND RESTRICTIONS

Apply only during alternate years in fields adjacent to aquatic areas such as lakes, reservoirs, rivers, permanent streams, marshes or natural ponds and estuaries. This product must not be applied within 150 feet (for aerial and air-blast applications), or 25 feet (for ground applications) from the aquatic areas listed above unless there is an untreated buffer area of that width between the area to be treated and the water body.

Do not use in greenhouses or other enclosed areas.

INTEGRATED PEST MANAGEMENT

Muscle ADV is a combination of systemic and contact fungicides with two different modes of action. Muscle ADV is an excellent disease control agent when used according to label directions for control of a broad spectrum of plant diseases. Muscle ADV is recommended for use in programs that are compatible with the principles of Integrated Pest Management (IPM), including the use of disease resistant crop varieties, cultural practices, pest scouting and disease forecasting systems which reduce unnecessary applications of pesticides.

FUNGICIDE RESISTANCE MANAGEMENT

Muscle ADV is effective for strategic use in programs that attempt to minimize disease resistance to fungicides. Some other fungicides which are at risk from disease resistance exhibit a single-site mode of fungicidal action. The chlorothalonil in Muscle ADV is a multi-site mode of action and may be used to delay or prevent the development of resistance to single-site fungicides. Consult with your federal or state Cooperative Extension Service representatives for guidance on the proper use of Muscle ADV in programs which seek to minimize the occurrence of disease resistance to other fungicides.

Limitations, Restrictions, and Exceptions

PEANUTS (SOILBORNE DISEASE)

Application Directions:

For control of soilborne diseases in a typical preventive fungicide program for peanuts, apply this product approximately 45 to 60 days after planting (early pegging). Repeat applications at 14-day intervals consecutively four times. Leaf spot advisory schedule: For control of soilborne diseases in an advisory schedule, apply Muscle ADV in the first advisory spray, usually occurring in July, and continue applications at 14-day intervals.

Do not feed hay or threshings to livestock, or allow livestock to graze in treated areas.

The active ingredient (tebuconazole) in this product must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by *Sclerotium rolfsii* and *Rhizoctonia solani*. Drought conditions may decrease the effectiveness of this product against root and pod rots.

Use this product in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices.

Do not apply more than 7.7 pints of this product (equal to 13 oz of tebuconazole ai/A and 2.9 lbs chlorothalonil ai/A) per acre per season.

Method

[Spray](#)

Pre-Harvest Interval

14 days

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[Postplant](#)