

FOLIAR APPLICATION - TREE

General Information

PRODUCT INFORMATION

HUMIPLUS ULTIMATE Liquid Nutrients can be applied to most vegetable crops, row crops, deciduous fruit and nut trees, citrus, avocados, grapes, melons, ornamentals, turf, pasture, range grasses, and most other crops.

HUMIPLUS ULTIMATE is a new liquid plant food developed for use as a foliar feed, a regular plant food applied to the soil, and as a starter plant food with the seed or transplant.

HUMIPLUS ULTIMATE liquid nutrients contain ENHANCE' (THA) Technical Humic acids. ULTIMATE nutrients are beneficial in combination with plant food and non-phytotoxic when used as directed.

ULTIMATE nutrients with (THA) Technical Humic Acids are unique as they can be used in most all forms of liquid fertilizers. ULTIMATE nutrients can be banded at planting time, side-dressed or sprayed in water solutions directly on deficient plants. Humic Acids may aid in the uptake of micronutrients.

HUMIPLUS ULTIMATE liquid nutrients have corrected deficiencies of a great many row crops, vegetables, and ornamental plants under soil conditions ranging from high organic matter (muck) to very low organic matter and from strong acid soils (pH 3) to high alkaline soils (pH 8.5) containing considerable calcium carbonate (free lime). HUMIPLUS ULTIMATE is effective under dry land and irrigation farming conditions. Differences in soil conditions, climate and plant varieties will determine how much more effective HUMIPLUS ULTIMATE liquid nutrients are than other sources of nutrients.

Remember, these humic acid products are used to fortify, produce, set and hold. It is important that timely applications are made to achieve these results.

DILUTION RATES

Aircraft and low volume sprayer: Use a minimum of 5 gallons (19 liters) of water per

acre.

Conventional Sprayers: Use a minimum of 20 gallons (76 liters) of water per acre.

Dilute Spray: Use 200-500 gallons (757 - 1892 liters) of spray solution per acre.

Concentrated Spray: Use 50-150 gallons (189 - 568 liters) of concentrated spray solution per acre.

Note: A wetting agent or spreader can be used when applying HUMIPLUS ULTIMATE.

Limitations, Restrictions, and Exceptions

FOLIAR APPLICATION - TREE

- Start as early as possible, usually at bud stage or just prior to first bloom and continue at 10-15 day intervals. The second, third, and fourth applications should come at petal fall and then two more applications after petal fall. These timely applications will insure sufficient levels of nitrogen and phosphorous to hold the set intact.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Directed](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Timings

[At bud stage or just prior to first bloom.](#)