

WALNUTS AND ALMONDS

General Information

PRODUCT INFORMATION

HUMIPLUS STIMULENT Liquid Nutrients can be applied to most vegetable crops, row crops, deciduous fruit and nut trees, citrus, avocados, grapes, melons, ornamentals, turf, pasture, range grasses, and most other crops.

HUMIPLUS STIMULENT is a new liquid plant food developed for use as a foliar feed, a regular plant food applied to the soil, and as a starter plant food with the seed or transplant. HUMIPLUS STIMULENT liquid nutrients contain ENHANCE' (THA) Technical Humic acids. HUMIPLUS nutrients are beneficial in combination with plant food and nonphytotoxic when used as directed.

HUMIPLUS STIMULENT nutrients with (THA) Technical Humic Acids are unique as they can be used in most all forms of liquid fertilizers. HUMIPLUS nutrients can be banded at planting time, side-dressed or sprayed in water solutions directly on deficient plants. Humic Acids may aid in the uptake of micro-nutrients.

HUMIPLUS STIMULENT liquid nutrients have corrected deficiencies of a great many row crops, vegetables, and ornamental plants under soil conditions ranging from high organic matter (muck) to very low organic matter and from strong acid soils (pH 3) to high alkaline soils (pH 8.5) containing considerable calcium carbonate (free lime). HUMIPLUS can be effective under dry land and irrigation farming conditions. Differences in soil conditions, climate and plant varieties will determine how much more effective HUMIPLUS liquid nutrients are than other sources of nutrients.

Remember, these humic acid products are used to fortify, produce, set and hold. It is important that timely applications are made to achieve these results.

Aircraft and low volume sprayers: Use a minimum of 10 gallons of water per acre.

Conventional sprayers: Use a minimum of 20 gallons of water per acre.

Limitations, Restrictions, and Exceptions

WALNUTS AND ALMONDS

- Apply three times each season starting at first full leaf.

Method

[Broadcast/Foliar Air](#)

[Directed](#)

[Band](#)

Rates

[field_rates 0](#)

•

Timings

[At first full leaf.](#)