

CORN (FIELD CORN, POPCORN) - PREPLANT, PREEMERGENCE

General Information

Product Information

HAVOC LV-Six herbicide is intended for selective control of many broadleaf weeds in certain crops, including, cereal grains (wheat, barley, millet, oats and rye), corn (field corn, popcorn and sweet corn), fallow land and crop stubble, potatoes, sorghum (grain and forage sorghum), and soybeans (preplant burndown application only), forests, rangeland and established grass pastures including Conservation Reserve Program (CRP) acres, non-cropland, grasses grown for seed or sod, and ornamental turf.

Apply HAVOC LV-Six as a water or oil-water spray during warm weather when weeds or woody plants are actively growing. Application under drought conditions will often give poor results. Use low spray pressure to minimize drift. Generally, the lower dosages specified on the label will be satisfactory for young, succulent growth of susceptible weed species. For less susceptible species and under conditions where control is more difficult, use higher specified rates. Deep-rooted perennial weeds such as Canada thistle and field bindweed and many woody plants usually require repeated applications for satisfactory control. Consult your State Agricultural Experiment stations or Extension Service Weed Specialists for recommendations based on use instructions of the label that best fit local conditions.

Use Precautions and Restrictions

Chemigation: Do not apply this product through any type of irrigation system.

Excessive amounts of 2,4-D in the soil may temporarily inhibit seed germination and plant growth.

Application Instructions

Spray Volume: Apply with calibrated air or ground equipment using sufficient spray volume to provide adequate coverage of target weeds or as otherwise directed in specific use directions. For broadcast application, apply the specified rate of this product in a spray volume of 2 or more gallons per acre by air and 10 or more

gallons per acre for ground equipment. Use low-pressure sprays to minimize drift. Where states have regulations, that specify minimum spray volumes, they should be observed. In general, spray volume should be increased as crop canopy, height and weed density increase in order to obtain adequate spray coverage. Do not apply less than 2 gallons total spray volume per acre.

Application Rates: Generally, lower rates in specified rate ranges will be satisfactory for more sensitive weeds species, when weeds are small, and when environmental conditions are favorable for rapid growth. Use higher rates in the specified rate range for less sensitive species and under less favorable growing conditions. For crop uses, do not mix with oil or other adjuvants unless specifically recommended on the label. Deep-rooted perennial weeds such as Canada thistle and field bindweed and many woody plants usually require repeated applications for effective control.

Spot Treatments

To prevent misapplication, spot treatments should be applied with a calibrated boom or with hand sprayers using a fixed spray volume per 1,000 sq ft as indicated below.

Hand-Held Sprayers: Hand-held sprayers may be used for spot applications of HAVOC LV-Six. Care should be taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based on the application rate for an area of 1,000 sq ft. Mix the amount of HAVOC LV-Six (fl oz or ml) corresponding to the desired broadcast rate in 1 to 3 gallons of spray. To calculate the amount of HAVOC LV-Six required for larger areas, multiply the table value (fl oz or ml) by the thousands of sq ft to be treated. An area of 1000 sq ft is approximately 10.5 X 10.5 yards (strides) in size.

Limitations, Restrictions, and Exceptions

Corn (Field Corn, Popcorn and Sweet Corn)

Specific Use Directions

For best results, growth conditions should be favorable for active weed growth. Use high rate in rate range for less susceptible weeds, cover crops such as alfalfa, weeds in advanced stages of development, or under less favorable growth

conditions.

Preplant: Apply 7 to 14 days before planting corn to control emerged broadleaf weed seedlings or existing cover crops.

Preemergence: Apply any time after planting, but before corn emerges to control broadleaf weed seedlings or existing cover crops. Do not use on light sandy soils.

Restrictions (Field Corn and Popcorn):

- Use 2 or more gallons of spray solution per acre.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

7 days

Rates

[field_rates 0](#)

•

Timings

[Preemergence \(Crop\)](#)

[Preplant](#)