

ANNUAL AND PERENNIAL GRASSES GROWN FOR SEED AND HAY, AND CONSERVATION RESERVE PROGRAM ACRES (CRP) - WEEDS PARTIALLY CONTROLLED

General Information

USE INFORMATION

Huskie Herbicide is a selective postemergence herbicide for control of important broadleaf weeds in spring, durum, winter wheat, barley, CRP, grasses grown for seed, rye, grain sorghum (to include grain and forage), and triticale.

ENVIRONMENTAL AND BIOLOGICAL ACTIVITY

Huskie Herbicide is a postemergence herbicide and best results are obtained when applications are made to young actively growing broadleaf weeds. Huskie Herbicide is primarily absorbed through the foliage and rapidly inhibits photosynthesis and pigment synthesis, causing death in susceptible weeds. Thorough spray coverage is important.

FIELD BIOASSAY

A field bioassay must be conducted for crops not listed on the label. To conduct a field bioassay, plant strips of the crop you want to grow the season following Huskie Herbicide application. Monitor the crop for response to Huskie Herbicide to determine if the crop can be grown safely in previously treated Huskie Herbicide areas.

Do not plant any rotational crop within 30 days following a Huskie Herbicide application.

WEED RESISTANCE

Huskie Herbicide contains active ingredients which inhibit photosynthesis and the HPPD enzyme systems which are members of WSSA Group 6 and 27 in susceptible plants. Huskie Herbicide may be an effective tool in the management of weed populations containing resistance to ALS, phenoxy or glyphosate herbicide modes of action. Repeated use of herbicides with the same mode of action allows resistant

weeds to spread. To manage the spread of resistant weed populations, use herbicides with different modes of action in tank mixture, rotation, or in conjunction with alternate cultural practices.

APPLICATION METHODS

Ground Application

Properly calibrated ground application equipment may be used to apply Huskie Herbicide postemergence as a foliar spray. Select spray nozzles that provide best spray distribution and weed coverage at the appropriate spray pressure. Avoid uneven spray distribution, skips, overlaps, and spray drift.

Apply 11 - 15 fl oz/A of Huskie Herbicide to labeled crops from fully expanded first true leaf up to flag leaf emergence. For most consistent control or under adverse growing conditions add AMS or an ammonium nitrogen source as directed under SPRAY ADDITIVES section. Do not use less than 11 fl oz/A of Huskie Herbicide unless directed by a Bayer CropScience representative.

Apply the appropriate dosage broadcast in 10 or more gallons of water per acre.

Use nozzles and spray pressure for ground application that deliver medium spray droplets as indicated in the nozzle manufacturer's catalogs such as 80-degree or 110-degree flat-fan nozzles in accordance with ASAE Standard S-572 for optimum spray coverage and canopy penetration. Use screens that are 50 mesh or larger.

Do not use flood-jet nozzles or cone nozzles. Nozzle types, nozzle spacings and lower spray pressures that produce coarse spray droplets may not provide adequate coverage of the weeds to ensure optimum control.

See the Spray Drift Management section of the label for additional information on proper application of Huskie Herbicide.

See Supplemental label for application of Huskie Herbicide in tank mixture with Osprey, Orion and Starane Herbicide for use and distribution only in the states of Washington, Oregon, Idaho and Washington.

Refer to the Supplemental label for application of Huskie Herbicide in tank mixture with MCPA Ester or Buctril 4 EC for use in Wheat or Barley in the States of Colorado and Idaho.

Refer to the Supplemental label for application of Huskie Herbicide in tank mixture with Osprey Herbicide and Starane Flex for broadleaf control in winter wheat in the States of Idaho, Oregon and Washington.

Limitations, Restrictions, and Exceptions

ANNUAL AND PERENNIAL GRASSES GROWN FOR SEED AND HAY, AND CONSERVATION RESERVE PROGRAM ACRES (CRP)

Huskie Herbicide may be applied to conservation reserve acres and certain annual and established perennial grasses grown for seed and hay for the management of broadleaf weeds. This product is not for use in sod production.

CROPS AND STAGE OF GROWTH AT APPLICATION

Huskie Herbicide may be applied from preemergence to established perennial ryegrass, annual ryegrass, tall fescue, fine fescue, Kentucky bluegrass and orchardgrass. Huskie Herbicide may be applied to established timothy.

HUSKIE HERBICIDE APPLICATION RATE

Apply 13.5 - 15 ounces of Huskie Herbicide per application per acre depending on the target weed species. Two applications of Huskie Herbicide can be made per year separated by at least 30 days.

APPLICATION METHODS

Ground Application (ONLY)

Properly calibrated ground application equipment may be used to apply Huskie Herbicide postemergence as a foliar spray. Select spray nozzles that provide best spray distribution and weed coverage at the appropriate spray pressure. Avoid uneven spray distribution, skips, overlaps, and spray drift.

Apply the appropriate dosage of Huskie Herbicide broadcast in 10 or more gallons of water per acre to labeled crops listed in the CROPS AND STAGE OF GROWTH AT

APPLICATION section of the label.

Under conditions where large weeds or dense weed populations are present or adverse environmental conditions exist, a greater spray volume of 15 – 20 gallons of spray solution per acre is required for best weed control. Do not apply with hollow cone type nozzles or other nozzles that produce a fine droplet spray. Use nozzles and spray pressure for ground application that deliver medium spray droplets as indicated in the nozzle manufacturer's catalogs such as 80-degree or 110-degree flat-fan nozzles in accordance with ASABE Standard S-572.1 for optimum spray coverage and canopy penetration. Use screens that are 50 mesh or larger.

Do not use flood-jet nozzles or cone nozzles. Nozzle types, nozzle spacings and lower spray pressures that produce coarse spray droplets may not provide adequate coverage of the weeds to ensure optimum control.

See the Spray Drift Management section of the label for additional information on proper application of Huskie.

Ground Application Restrictions:

- Do not apply this product with backpack or hand-held application equipment.

WEED CONTROL IN GRASSES GROWN FOR SEED, HAY, AND CRP

Apply Huskie Herbicide as directed to control many important broadleaf weeds in labeled grasses grown for seed and hay.

RESTRICTIONS FOR HUSKIE HERBICIDE USE IN CRP and GRASSES GROWN FOR SEED AND HAY:

- Do not apply more than two applications of Huskie Herbicide per acre per year.
- Grass forage may be cut or grazed seven days after application but do not cut for hay within 30 days after treatment.
- Aerial and chemigation application are prohibited.

NOTES:

- Chickweed, common; Horseweed/Marestail; Mayweed chamomile / dogfennel:
Partially controlled weeds may be stunted in growth and/or be reduced populations
as compared to non-treated areas but control will generally not be commercially
acceptable.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[Preemergence \(Crop\)](#)

[Postemergence \(Weed\)](#)