

LANDSCAPE ORNAMENTALS - ANNUAL BROADLEAF WEEDS

General Information

GENERAL INFORMATION

This product is a preemergence herbicide for control of certain annual grasses and broadleaf weeds in container grown ornamentals, landscape ornamentals, nursery stock, ground covers, established flowers, ornamental bulbs, non-bearing fruit and nut trees and non-bearing vineyards. Apply this product before or after planting but prior to germination of target weeds, or immediately after cultivation. Length of weed control will vary with weed population, potting media or soil conditions, temperature, watering regime, and other factors. Following application, user should monitor and observe level of weed control over time to determine when additional applications may be needed. Repeat applications should not be made sooner than 60 days after a previous application of this product. Do not apply over 240 pounds per acre total of this product within a 12 month period.

This product does not control established weeds. Existing weeds should be controlled by cultivation or with postemergence herbicides. Weed residues, prunings and trash should be removed or thoroughly mixed into soil prior to treatment. Soil should be in good condition and free of clods at the time of application. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation, is required to activate this product. Optimum weed control is obtained when this product is activated within 3 days of application. If rainfall or irrigation has not occurred within 3 days of application and tillage is possible, this product may be activated using cultivation equipment capable of uniformly mixing the herbicide into the upper 1-2 inches of soil. Failure to activate this product within 3 days of application may result in erratic weed control. Do not apply when wind conditions favor drift of this product's granules from the target area.

Optimum weed control will be obtained when followed by overhead irrigation or rainfall within a few hours after surface application.

GENERAL PRECAUTIONS AND RESTRICTIONS

Entry Restrictions: Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the

incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

To avoid possible injury, do not apply this product to:

- cuttings that have been planted in post for the first time
- * ground covers until they are established and well rooted

Do not apply this product to newly transplanted ornamentals, ground covers, flowers and non-bearing fruit and nut trees until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or injury may occur.

Do not make preplant applications of this product to areas where gladioli corns less than one inch in diameter will be planted or injury may occur.

Do not apply this product in greenhouses or other enclosed structures.

Repeat applications should not be made sooner than 60 days after a previous application of this product. Do not apply over 240 pounds per acre total of this product within a 12 month period.

Users who wish to use this product on plant species not recommended on this label may determine suitability for such uses by making trial application of this product at a recommended rate to small number of plants. Prior to using this product on a larger number of plants, the treated plants should be observed for signs of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. The user assumes responsibility for any plant damage or other liability resulting from the use of this product on plant species not recommended on this label.

APPLICATION DIRECTIONS

Apply this product using a drop or rotary-type spreader designed to apply granular herbicides or insecticides. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working

properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or plant injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

APPLICATION TECHNIQUES FOR APPLYING THIS PRODUCT

When using a drop-type spreader, a splash board mounted under the hopper will provide more even granule distribution.

A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.

For treating smaller areas or rows of nursery stock or ornamental beds, a hand-held or push-type rotary applicator such as whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.

A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, hard to treat areas. Carefully measure the amount of product needed to avoid over-application.

Limitations, Restrictions, and Exceptions

Landscape Ornamentals, Ground Covers, Established Flowers, Ornamental Bulbs

This product is recommended as a preemergence treatment for control of certain annual grasses and broadleaf weeds in landscape ornamentals, ground covers, established flowers, ornamental bulbs. Apply this product before or after planting but prior to germination of target weeds, or immediately after cultivation.

Refer to the "General Information", "General Precaution and Restrictions" and "Application Directions" sections of the label for general use precautions and information on application, application rates and weeds controlled.

- In addition to the weeds controlled, the following weeds will be partially controlled or suppressed

ORNAMENTAL BULBS:

This product may be applied for control of susceptible annual weeds in ornamental bulbs, e.g. bulbous iris, daffodil (narcissus), hyacinth and tulip. Apply this product to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual weeds. This product may also be applied following bulb emergence. For fall planted bulbs, apply this product again in late winter or early spring to weed-free soil surfaces.

- Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants which will not bear fruit for at least one year after treatment.

Method

[Surface](#)

[Irrigation](#)

[Surface](#)

[Irrigation](#)

[Surface](#)

[Irrigation](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Timings

[Preplant](#)

[Preemergence \(Weed\)](#)

[Postplant](#)