

RED CLOVER, ALFALFA - EARLY BUD STAGE FOR POWDERY MILDEW

General Information

INSTRUCTIONS FOR USE

Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality.

USE PRECAUTIONS: Do not use this product within 30 days of an oil spray at any stage other than dormant (deciduous only) unless prior experience in your immediate area has shown that shorter intervals will not result in phytotoxic injury. Lime-Sulfur may burn foliage when temperature is high; avoid applying this material during very hot periods of the day (above 85°F). Keep container tightly closed when not in use. If crust should form on the surface, break through the crust so product will pour.

RESTRICTIONS: Not for residential use or application to residential sites. This product may not be used in, on, or around any structure, vehicle, article, surface or area associated with the household, including non-agricultural outbuildings, non-commercial greenhouses, pleasure boats and recreational vehicles; in or around any preschool or day care facility or on humans or pets.

Do not apply, or allow to drift, to painted surfaces, as a permanent stain may result. Do not combine Lime-Sulfur with other pesticides unless previous experience has shown them to be compatible. If tank mixing, always follow the more restrictive label directions. This product cannot be mixed with any product containing a label prohibition against such mixing. Read entire label and use strictly in accordance with label directions.

Do not apply this product through any type of irrigation system.

Do not use this product for crop thinning.

Unless otherwise indicated, the following use rates are per 100 gallons of water. Where a rate range is given, use the higher rates when disease is severe or where

disease was severe in the previous season.

Do not apply more than 30 gallons of this product/acre/year to any crop listed on the label.

Do not apply this product to harvested fruit.

DILUTE SPRAY OR APPLICATION - For general dilute applications, use higher gallonage (100 - 400 gal/acre) on larger trees where required.

CONCENTRATE SPRAY OR APPLICATION - For concentrate spray applications, use 100 gal/acre.

Restriction: When using concentrate sprays do not exceed 12 gallons of LIME-SULFUR SOLUTION per 100 gallons on a per acre basis. (Rust mites and scale insects have been shown to be more effectively controlled with dilute or semi-dilute applications.)

Limitations, Restrictions, and Exceptions

RED CLOVER, ALFALFA

- Apply 1 gallon per acre in 50 to 100 gallons of spray.

Method

[Spray](#)

Rates

[field rates 0](#)

•

Restricted Entry Interval

48 hours

Timings

[At first sign of disease.](#)

[Early bud stage.](#)