

ANIMAL REPELLANCY APPLICATIONS (APPLICATIONS TO VINES, TREE TRUNKS AND DORMANT PLANTS)

General Information

GENERAL PRODUCT INFORMATION

Thiram Granuflo Agricultural Fungicide is a broad-spectrum fungicide for plant disease control in the culture of Peaches and Strawberries. Thiram Granuflo Agricultural Fungicide is a broad-spectrum fungicide for use on Turf (golf courses) to control Dollar Spot, Brown Patch and Fusarium Patch. When used as a Seed Protectant, Thiram Granuflo Agricultural Fungicide will usually increase crop stands and yields by reducing seedling losses from seed decay, damping-off or seedling blights. When used as a Repellent, Thiram Granuflo Agricultural Fungicide can protect trees, vines, nursery stock, shrubs and ornamentals from rabbit, deer and rodent depredation.

Taminco Inc.'s recommendations for the use of this product are based upon tests believed to be reliable. FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THE LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

Not for use on turf being grown for sale or other commercial use such as sod, or for commercial seed production, or for research purposes.

Do not use on residential lawns or turf.

GENERAL RECOMMENDATIONS

Be sure spray equipment is clean before use. If it is contaminated with any material, crop injury or sprayer clogging may result. Calibrate spray equipment before use.

For agricultural foliar applications use recommended rates for dilute, concentrate and aerial applications. Ground application is preferred, especially when full canopy of foliage exists. For dilute application, use Thiram Granuflo Agricultural Fungicide at the recommended rate listed in the "per 100 gallons" column of the Specific Applications table. For concentrate ground application, apply the recommended rate listed in the "per acre" column of the Specific Applications table in a minimum

of 20 gallons per acre. For aerial application, apply the recommended rate listed in the \"per acre\" column of the Specific Applications table in a minimum of 10 gallons per acre. Applications of Thiram Granuflo Agricultural Fungicide are most effective when made before infection periods. Under low disease conditions, minimum label rates should be used; however, maximum label rates and shortened spray schedules are recommended for severe or threatening disease conditions. If you are not familiar with such conditions, get advice from your state agricultural extension service.

IMPORTANT: When spraying, thorough coverage is necessary to provide good disease control. Addition of a spreader/sticker to the finished spray mixture may improve coverage.

Make fresh spray mixtures for each application.

For sale to and use by professional applicators only. Not for sale to or use by homeowners/consumers.

Limitations, Restrictions, and Exceptions

ANIMAL REPELLANCY APPLICATIONS:

When used according to label directions, Thiram Granuflo Agricultural Fungicide can protect trees, vines, nursery stock, shrubs and ornamentals from rabbit, deer and rodent depredation. One application will protect treated parts from 1 to 3 months, depending upon weather conditions. Thiram Granuflo Agricultural Fungicide can also be used as a repellent in deterring deer from moving into important areas such as airport runways as a result of local feeding and browsing. It can also deter migratory birds such as geese on golf courses and other turf areas. The addition of a latex based sticker prolongs the residual effectiveness of the treatment.

Repellency is the result of rendering the treated plant parts distasteful. The maximum application rate for products applied as a mammal (deer) repellent is one pound of product in 3 - 7 gallons of water applied in an area of 1000 square feet. Applications to ornamentals will be restricted to cold weather areas, climate zones 1 through 7, in the following states (MA, NH, VT, ME, RI, CT, NY, NJ, PA, MD, DE, WV, OH, KY, MI, IN, IL, WI, MN, IA, MO, ND, SD, NE, KS, MT, WY, CO, ID, UT, WA, OR, NV, and AK) during the winter season only (October through March).

Applications To Vines, Tree Trunks and Dormant Plants:

Prepare a slurry of 1.33 pounds of Thiram Granuflo Agricultural Fungicide in 3 gallons of water and apply as a full coverage spray to all portions of the plant to be protected. For greater retention and residual efficacy, add 1 pint of a latex based sticker to the spray solution.

Method

[Spray](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

24 hours

Exception: Once treated seeds are planted in soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction, if there will be no contact with the treated seeds.

Timings

[N.A.](#)