

TURFGRASS - LEAF SPOT - CURATIVE APPLICATIONS

General Information

GENERAL INFORMATION

Mixing Procedures

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Agitation is necessary for proper dispersal of the product. Maintain agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

ARMADA 50 WDG Alone: Add 1/2 of the required water to the mix tank. With the agitator running, add the prescribed dose of ARMADA 50 WDG to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after ARMADA 50 WDG has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Resistance Management: ARMADA 50 WDG contains the strobilurin class of chemistry, which exhibits no known cross-resistance to other chemical classes including sterol inhibitors, dicarboximides, benzimidazoles, anilinopyrimidines, or phenylamides. However, certain fungal pathogens are known to develop resistance to products used repeatedly. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies. Such strategies may include rotating and/or tank mixing with products having different modes of action; or limiting the total number of applications per season. Bayer encourages responsible product stewardship to ensure effective long-term control of the fungal diseases on the label. See specific recommendations in the turf section and the ornamentals section.

PLANT HEALTH

Apply ARMADA 50 WDG at 1.2 to 1.5 oz/1,000 sq ft to sod production of warm season grasses such as St. Augustine's, centipede, and Bermuda grass. The following plant health enhancements will be seen 30 to 60 days after application of

ARMADA 50 WDG; visible leaf greenness and density. Sod laid under shade trees also benefits in turf quality when this application is made prior to transfer. Apply ARMADA 50 WDG at 1.2-1.5 oz /1,000 sq ft 15-30 days pre-harvest for plant health benefits.

Refer to label for more informations.

Limitations, Restrictions, and Exceptions

TURFGRASS - LEAF SPOT - CURATIVE APPLICATIONS

Application Interval/Timing: 21-28 days.

For preventive applications apply when conditions are favorable for disease. Use the higher rate/shorter interval for curative applications or under severe disease conditions.

Method

[Spray](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[N. A.](#)