

## **FRUITS - PEAR - APHID**

### General Information

### DIRECTIONS FOR USE

#### When using Ortho Dial 'N Spray

1. Set dial to the setting indicated in the use directions for your specific application.
2. Pour product into sprayer to fill jar one-quarter to one-half full.
3. After spraying, unused product must be poured back into its original container.

#### When using a tank sprayer

1. Blend appropriate amount of Ortho MAX Malathion Insect Spray Concentrate with appropriate amount of water following the dilution instructions for each site-pest combination listed.
2. Operate sprayer by following the manufacturer's instructions. Spray as directed.

### Limitations, Restrictions, and Exceptions

### FRUITS

- Malathion may cause injury to Bosc Pears and Ribier Grapes.

### USE THIS AMOUNT

- Ortho Dial 'N Spray setting is 2 tsp.

### HOW TO APPLY

- Use an Ortho Dial 'N Spray or tank sprayer.
- Spray thoroughly covering both upper and lower leaf surfaces and other infested plant parts.

### WHEN TO APPLY

- Apply as necessary to control listed insects; waiting at least 30 days between each

application.

Method

[Spray](#)

Pre-Harvest Interval

1 day

Rates

[field\\_rates 0](#)

•

Timings

[N.A.](#)