

FRUITS - CHERRIES, GRAPES & STRAWBERRIES - APHID

General Information

DIRECTIONS FOR USE

When using Ortho Dial 'N Spray

1. Set dial to the setting indicated in the use directions for your specific application.
2. Pour product into sprayer to fill jar one-quarter to one-half full.
3. After spraying, unused product must be poured back into its original container.

When using a tank sprayer

1. Blend appropriate amount of Ortho MAX Malathion Insect Spray Concentrate with appropriate amount of water following the dilution instructions for each site-pest combination listed.
2. Operate sprayer by following the manufacturer's instructions. Spray as directed.

Limitations, Restrictions, and Exceptions

FRUITS

- Malathion may cause injury to Bosc Pears and Ribier Grapes.

USE THIS AMOUNT

- Ortho Dial 'N Spray setting is 2 tsp.

HOW TO APPLY

- Use an Ortho Dial 'N Spray or tank sprayer.
- Spray thoroughly covering both upper and lower leaf surfaces and other infested plant parts.

WHEN TO APPLY

- Apply as necessary to control listed insects; waiting at least 30 days between each

application.

Method

[Spray](#)

Pre-Harvest Interval

3 days

Rates

[field_rates 0](#)

•

Timings

[N.A.](#)