

LAWN - CRANE FLIES

General Information

DIRECTIONS FOR USE

with its labeling.

Do not apply a broadcast application to interior surfaces of homes.

Do not apply by air.

Do not apply in greenhouses, nurseries.

Do not apply this product through any kind of irrigation system.

Not for use on sod farm turf, golf course turf, or grass grown for seed.

Do not water treated area to the point of run-off.

Do not make applications during rain.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

Additional Application Restrictions for Residential Outdoor Surface and Space Sprays:

All outdoor applications, if permitted elsewhere on the label, must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses, if allowed elsewhere on the label:

1. Applications to soil or vegetation, as listed on the label, around structures;
2. Applications to lawns, turf, and other vegetation, as listed on the label;
3. Applications to the side of a building, up to a maximum height of 3 feet above grade;
4. Applications to underside of eaves, soffits, doors, or windows permanently

protected from rainfall by a covering, overhang, awning, or other structure;

5. Applications around potential pest entry points into buildings, when limited to a surface band not to exceed one inch in width;

6. Applications made through the use of a coarse, low pressure spray to only those portions of surfaces that are directly above bare soil, lawn, turf, mulch or other vegetation, as listed on the label, and not over an impervious surface, drainage or other condition that could result in runoff into storm drains, drainage ditches, gutters or surface waters, in order to control occasional invaders or aggregating pests.

General Information on the Use of this Product

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended only for aesthetic purposes or climatic modifications and being grown in interior plantscapes, ornamental gardens or parks, or lawns and grounds.

General Application Instructions

Ortho MAX PRO formulation mixes readily with water and other aqueous carriers, and controls a wide spectrum of insects and mites on trees, shrubs, foliage plants, non-bearing fruit and nut trees, and flowers in interiorscapes including hotels, shopping malls, office buildings, etc., and, outdoor plantscapes, such as around residential dwellings, parks, institutional buildings, recreational areas, athletic fields and home lawns. Nonbearing crops are perennial crops that will not produce a harvestable raw agricultural commodity during the season of application.

Ortho MAX PRO may be tank-mixed with other pesticides, including insect growth regulators. When tank mixing Ortho MAX PRO with other pesticides, observe all precautions and limitations on each separate product label. The physical compatibility of Ortho MAX PRO may vary with different sources of pesticide products, and local cultural practices. Any tank mixture which has not been previously tested should be prepared on a small scale (pint or quart jar), using the proper proportions of pesticides and water to ensure the physical compatibility of the mixture.

The following procedure is recommended for preparation of a new tank mix, unless

specified otherwise in label directions: (1) Add wettable powders to tank water, (2) Agitate, (3) Add liquids and flowables, (4) Agitate, (5) Add emulsifiable concentrates, and (6) Agitate. If a mixture is found to be incompatible following this order of addition, try reversing the order of addition, or increase the volume of water. Note: If the tank-mixture is found to be compatible after increasing the amount of water, then the sprayer will need to be recalibrated for a higher volume application. Do not allow tank mix to stand overnight.

Resistance: Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state pest management authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and suspect that resistance is a reasonable cause, immediately consult your local company representative or pest management advisor for the best alternative method of control for your area.

- Refer in the label for Dilution Chart.

Foam Applications

Ortho MAX PRO may be converted to foam and used to treat void spaces, or as a spot spray on vertical or horizontal surfaces where visual marking of application is desired. Use of a foaming agent increases a.i. surface contact time on challenging surfaces and provides visual marking of the application. Ensure that the foaming agent is approved for food surface/area contact use.

Attention

Do not apply a broadcast application to interior surfaces of homes.

Do not apply to pets, crops, or sources of electricity.

Firewood is not to be burned for one month after treatment.

Use only in well ventilated areas.

During any application to overhead areas within the structure, cover surfaces below with plastic sheeting or similar material, except for soil surfaces in crawlspaces.

Do not allow spray to contact food, foodstuffs, food contacting surfaces, food utensils or water supplies.

Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product.

Do not treat areas where food is exposed.

During indoor surface applications do not allow dripping or run-off to occur.

Do not allow people or pets on treated surfaces until spray has dried.

Let surfaces dry before allowing people and pets to contact surfaces.

Prior to applying Ortho MAX PRO to wood siding, especially rough wood siding, be sure to thoroughly agitate the tank mixture. Prior to treating wood siding, test a small area and allow it to dry to be sure no deposits will form. Follow the same procedure when applying to wood surfaces in direct sunlight or the heat of the day.

Do not apply this product in patient rooms or in any rooms while occupied by the elderly or infirm.

Do not apply in classrooms when in use.

Do not apply when occupants are present in the immediate area in institutions such as libraries, sports facilities, etc.

Application equipment that delivers low volume treatments, such as the Micro-Injector or Actisol applicators, may also be used to make crack and crevice, deep harborage, spot and general surface treatments of Ortho MAX PRO.

Limitations, Restrictions, and Exceptions

APPLICATION RECOMMENDATIONS

LAWN: Apply Ortho MAX PRO as a broadcast treatment. Use application volumes of

up to 10 gallons per 1000 square feet to get uniform coverage when treating dense grass foliage.

For low volume applications, less than 2 gallons/1000 square feet, immediate irrigation of treated area with at least 0.25 inches of water following application to ensure efficacy of sub-surface pests such as, but not limited to, Mole Crickets, is recommended.

LAWN APPLICATION RATES

The application rates listed will provide excellent control of the respective pests under typical conditions. However, at the discretion of the applicator, Ortho MAX PRO may be applied at up to 1 fl. oz. per 1000 square feet to control each of the pests listed in this Table. The higher application rates should be used when maximum residual control is desired or heavy pest populations occur.

- In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

- In New York State, do make a single repeat application of Ortho MAX PRO if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

COMMENTS

Crane Flies: Treatments can be made to control early to mid-season larvae (approximately August - February) as they feed on plant crowns. Treatments made to late-season larvae (approximately March, April) may only provide suppression.

Method

[Spray](#)

Rates

[field_rates 0](#)

•

Timings

[Crane Flies: Early to mid-season larvae \(approximately August - February\) as they feed on plant crowns.](#)