

CITRUS (CA) - MITES AND SCALES (CENTRAL CALIFORNIA)

General Information

PRECAUTIONS AND RESTRICTIONS

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Only protected handlers may be in the area during application.
- Do not apply this product through any type of irrigation system.
- Do not apply if this product will drift to food forage or other crops that might be damaged.
- For applications, the spray tank must be free of any sulfur residue before using PHT 440 Supreme Spray Oil. Do not combine this product with any type of sulfur spray. Do not apply PHT 440 Supreme Spray Oil after a sulfur application until sufficient time has passed for safety (at least 30 days). Do not apply sulfur after an application of PHT 440 Supreme Spray Oil until sufficient time has passed for safety (at least 30 days). If doubts exists, spray PHT 440 Supreme Spray Oil mixture on a portion of one tree that was treated with sulfur. If no burn is noted in several days, enough time has passed. Do not spray when temperatures are above 95° F or when temperatures drop below 35° F. Do not use on foliage closer than 60 days after or 90 days before using Captan or Folpet products.
- PHT 440 Supreme Spray Oil must be diluted before an application is made.
- DO NOT MIX with insecticides or fungicides incompatible with oil. Consult your State Agricultural Extension Service or Fieldman for specific compatibilities.
- Application to trees weakened by disease, drought, drying winds or high nitrogen applications may result in injury to the trees.

ORCHARD CONCENTRATE APPLICATION

Apply according to rates given in the table for each crop. All quantities of spray solution are for mature trees and vines with normal foliage cover. Reduce quantities

proportionally for smaller immature trees and vines. Use only on crops listed on the label. Special concentrate spray equipment is necessary for this application.

Restriction: Use efficient equipment of the proper type. Do not spray during or immediately prior to hot or freezing weather (over 95° or under 35°F), hot dry wind, rain or other unsuitable conditions avoid over spraying or double spraying. Spray plants only when in vigorous condition and when their moisture condition is suitable.

Limitations, Restrictions, and Exceptions

CITRUS

Central California: Apply 0.5 to 1.4 gallons per 100 gallons of water as thorough coverage spray (TC).

Alternately, use a low volume spray (LV) to apply 2.5 to 15 gallons per acre.

Do not apply to lemons within 60 days of harvest if lemons are to be sweated.

For lemons and limes, make applications in June through September. For grapefruits make applications in August through September. For oranges, make applications on navels and Valencias - July through August and on tangelos, tangerines, and other citrus varieties - July through August.

For navels and Valencias, treatment can be made from Feb 15 to 50% bloom, to avoid injury, use low concentration (1.0%).

On navel oranges, do not apply during or after color break.

Maximum of 212 pounds active ingredient per acre (maximum 2000 gallons of spray mix per acre) in California.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[Lemons and Limes: In June through September.](#)

[Grapefruits: In August through September.](#)

[Oranges \(Navels and Valencias\): July through August.](#)

[Tangelos, tangerines, and other citrus varieties: July through August.](#)