

NON-FOOD CONTACT PAPER USES: TREATMENT OF STARCH USED FOR SIZING ON THE PAPER MACHINE (SHOCK OR INTERMITTENT DOSING)

General Information

DIRECTIONS FOR USE

CONTROL OF ALGAL, FUNGAL, AND BACTERIAL GROWTH IN PULP AND PAPER MILL SYSTEMS FOR FOOD AND NON-FOOD CONTACT PAPER

Proxitane WW -12 provides an effective means to treat various process waters for slime control. Dosage rates should be increased or decreased depending on control achieved.

Maximum usage rate must not exceed 2 lbs Proxitane WW-12 solution per ton (2000 lbs., dry basis) of pulp or paper produced.

Limitations, Restrictions, and Exceptions

CONTROL OF ALGAL, FUNGAL, AND BACTERIAL GROWTH FOR NON-FOOD CONTACT PAPER USES.

TREATMENT OF STARCH USED FOR SIZING ON THE PAPER MACHINE - Apply Proxitane WW-12 directly to the starch storage tank or through the recirculation loop. Apply with either shock, intermittent, or continuous dosing. Shock doses may be applied for 1 to 2 hours, whereas intermittent doses may be applied for 5 to 60 minutes up to 12 times per day. For either shock or intermittent dosing, apply 0.8 to 5 gallons Proxitane WW -12 per 1000 gallons of starch solution to achieve 100 to 600 ppm 100% peracetic acid.

Method

[N. A.](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Timings

[N. A.](#)