

## **TURF APPLICATIONS (CURATIVE TREATMENT) - ALGAECIDE/BACTERICIDE TREATMENT**

General Information

### **DIRECTIONS FOR USE**

#### **CONTROL OF ALGAL, FUNGAL, AND BACTERIAL GROWTH IN PULP AND PAPER MILL SYSTEMS FOR FOOD AND NON-FOOD CONTACT PAPER**

Proxitane WW -12 provides an effective means to treat various process waters for slime control. Dosage rates should be increased or decreased depending on control achieved.

Maximum usage rate must not exceed 2 lbs Proxitane WW-12 solution per ton (2000 lbs., dry basis) of pulp or paper produced.

Limitations, Restrictions, and Exceptions

### **TURF APPLICATIONS**

Broad spectrum treatment for control of algae, fungi and bacteria on turf. For use on all turf types such as commercial turf, lawns, athletic fields and golf course fairways, greens and tees. Use Proxitane WW -12 to control Anthracnose, Brown Spot, Dollar Spot, Copper Spot, Fairy Ring, Pink Snow Mold, Pythium, Phytophthora, Summer Patch, Rhizoctonia, Scum, Fusarium, Blight, Stripe Smut, Leaf Spot, Algae, Slime Molds and their spores. Proxitane WW -12 controls on contact.

For algicide/bactericide treatment, use Proxitane® WW -12 to control algae and bacterial diseases and the odors and the conditions these organisms may cause.

1. Typical treatment rates involve treating approximately 1000 square feet of lawn area with 1 to 10 gallons of diluted solution. Spray entire area until run-off; saturation of the entire area will help ensure the solution penetrates algal crusts and deposits. Add a spreader surfactant for best results.
2. For initial (curative) treatment of heavy infestations of algae or bacterial disease, dilute 1.6 to 6.5 fl. oz. in 5 gallons of clean water. Apply 5-10 gallons of dilute

solution per 1000 square feet.

4. Repeat applications every 5 to 7 days or as needed to control new or established disease conditions. For best results, apply immediately after grass has been cut.

Optimum treatment time is early morning or late afternoon. Applications can be made during wet or rainy weather. Use spray solution the same day it is prepared; do not store and reuse mixed spray solution. Proxitane WW -12 can be injected through automatic irrigation systems in turf areas. Refer to Chemigation Directions for Use for specific instructions on using this product through irrigation systems.

Method

[Spray](#)

Rates

[field\\_rates 0](#)

•

Timings

[After grass has been cut.](#)