

TURF APPLICATIONS (CURATIVE TREATMENT) - FUNGICIDE TREATMENT

General Information

DIRECTIONS FOR USE

CONTROL OF ALGAL, FUNGAL, AND BACTERIAL GROWTH IN PULP AND PAPER MILL SYSTEMS FOR FOOD AND NON-FOOD CONTACT PAPER

Proxitane WW -12 provides an effective means to treat various process waters for slime control. Dosage rates should be increased or decreased depending on control achieved.

Maximum usage rate must not exceed 2 lbs Proxitane WW-12 solution per ton (2000 lbs., dry basis) of pulp or paper produced.

Limitations, Restrictions, and Exceptions

TURF APPLICATIONS

For fungicide treatment of turf, use on golf course fairways, greens and tees of Bentgrass, Bluegrass, Bermudagrass, Fescue, Ryegrass, St. Augustinegrass and their mixtures.

1. Typical treatment rates involve treating approximately 1000 square feet of lawn area with 1 to 10 gallons of diluted solution. Spray entire area until run-off. Add a spreader surfactant for best results.
2. Start applications at the first sign of disease and repeat every 5 to 7 days or as needed to control new or established disease conditions. For best results, apply uniformly over the area immediately after grass has been cut.
3. For initial (curative) treatment of heavy infestations of fungal disease, dilute 0.8 to 3.2 fl. oz. in 5 gallons of clean water. Apply 5 - 10 gallons of dilute solution per 1000 square feet.

Optimum treatment time is early morning or late afternoon. Applications can be made during wet or rainy weather. Use spray solution the same day it is prepared;

do not store mixed spray solution for later use.

Method

[Spray](#)

Rates

[field rates 0](#)

•

Timings

[After grass has been cut.](#)

[At the first sign of disease and repeat every 5 to 7 days or as needed to control new or established disease conditions.](#)