

TURF APPLICATIONS (SOIL TREATMENT FOLLOWING INOCULATION OF SOIL WITH BENEFICIAL MICROORGANISMS) - FUNGICIDE TREATMENT

General Information

DIRECTIONS FOR USE

CONTROL OF ALGAL, FUNGAL, AND BACTERIAL GROWTH IN PULP AND PAPER MILL SYSTEMS FOR FOOD AND NON-FOOD CONTACT PAPER

Proxitane WW -12 provides an effective means to treat various process waters for slime control. Dosage rates should be increased or decreased depending on control achieved.

Maximum usage rate must not exceed 2 lbs Proxitane WW-12 solution per ton (2000 lbs., dry basis) of pulp or paper produced.

Limitations, Restrictions, and Exceptions

TURF APPLICATIONS

To treat turf following inoculation of soil with beneficial microorganisms, use Proxitane WW -12 to control plant pathogens on the foliar portion of turf. Do not drench the root system, or a temporary reduction in beneficial soil microorganisms may occur. Use 1.1 to 2.2 fl. oz. per 5 gallons of clean water. Apply to the turf by lightly spraying leaf surfaces. Do not allow solution to be drenched into the soil and root systems. Drenching of Proxitane® WW -12 into the soil may result in temporary reduction of beneficial microorganisms.

Method

[Spray](#)

Rates

[field rates 0](#)

•

Timings

[N. A.](#)