

FRUIT DISEASES - PEACH, NECTARINE, APRICOT, ETC. - BROWN ROT BLOSSOM BLIGHT

General Information

GENERAL INFORMATION

CONTROLS: Leaf spots, rust, blights, fruit rots, mildews, scab, molds, etc.

USE ON: vegetables, fruit trees, flowers, shrubs, and shade trees

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, and grounds. Also for use on fruit trees and vegetables.

READ ENTIRE LABEL AND USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product is highly effective for the control of a broad spectrum of vegetable, fruit and ornamental plant diseases when it is used according to the directions on the label. When conditions are favorable for development, thorough uniform coverage of plant surfaces is essential for good disease control.

General Precautions and Restrictions: Do not allow people or pets to enter treated areas until sprays have dried. Do not apply this product in a way that will contact other persons or pets, either directly or through drift.

Limitations, Restrictions, and Exceptions

FRUIT DISEASES

Apply full coverage spray to the point of runoff to thoroughly cover tree canopy including undersides of leaves

PEACH, NECTARINE, APRICOT, CHERRY, PLUM, PRUNE: Brown Rot Blossom Blight - Make one application at popcorn (pink, red or early white bud stage when first color shows) and a second application at full bloom. If cool, moist weather continues, make an additional application at petal fall.

Method

[Spray](#)

Timings

[At popcorn \(pink, red or early white bud stage when first color shows\) and a second application at full bloom.](#)