

LANDSCAPE, ORNAMENTAL, AND TREE PESTS (OUTDOORS)

General Information

DIRECTIONS FOR USE

People and pets may enter treated area after application has dried.

This product has a long residual, providing up to 16 weeks control of cockroaches and up to 12 weeks of other pests.

FOR OUTDOOR USE ONLY.

DO NOT USE INSIDE DWELLINGS.

SHAKE PRODUCT WELL BEFORE USE.

- Virtually odor-free
- Fast-acting
- Easy-to-use
- Non-staining
- Kills Ants, Earwigs, Fleas, Insects, Mites, Roaches, Silverfish, Spiders, Ticks & Other Listed Pests

1 Tablespoon (1 Tbs.) = 3 teaspoons (tsp.);

1 fl. oz. = 2 Tbs.; 4 fl. oz. = ½ cup.

Do not use household utensils to measure this product.

Limitations, Restrictions, and Exceptions

LANDSCAPE, ORNAMENTAL, AND TREE PESTS (OUTDOORS)

HOW TO APPLY

- Spray evenly over foliage and treatment surfaces to the point of runoff.

WHEN TO REAPPLY

- Repeat application is necessary only if there are signs of renewed insect activity. Repeat application should be limited to no more than once per seven days.

MOSQUITOES (OUTDOORS)

Mosquitoes coming to rest on areas treated for the control of lawn and home-invading pests will be killed for varying periods of time after application, depending on exposure of treated areas to environmental conditions.

IMPORTANT

DO NOT treat structures that contain cisterns, wells, French drains or sump systems. Soil in the vicinity of wells, cisterns or ponds must not be treated. For those situations, contact your County Cooperative Extension Agent or consult a commercial pest control service. DO NOT treat soil that is water saturated or frozen, or in any conditions where run-off or movement from the treatment area (or site) is likely to occur. DO NOT apply while it is raining. DO NOT use on pets. DO NOT use on edible plants. Spray drift onto these plants must be rinsed off immediately with water.

Method

[Spray](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Timings

[For Beetles, Scale Crawlers, Twig Borers, and Weevils: Treat tree trunks, stems, and twigs in addition to plant foliage.](#)

[For Bagworms, apply when larvae begin to hatch and spray larvae directly. Application when larvae are young will be most effective.](#)