

## **USE IN KENNELS AND DOGHOUSES**

### General Information

### PRODUCT INFORMATION

PIVOT 10 inhibits the typical development and ability of immature insects from reaching adulthood. PIVOT 10 use can limit insect reproduction by causing adult insect sterilization, reduced egg production, behavior modifications, deformed insects and death. PIVOT 10 may be used in an Integrated Pest Management (IPM) program to help in the prevention of insect resistance.

### PRECAUTIONS:

1. Do not use this product in or on electrical equipment due to possibility of shock hazard.
2. Do not wet carpet, draperies and furniture. **STAINING:** Before using PIVOT 10, test the spray solution in a hidden area before use. As with other products, some fibers or surfaces (such as vinyl siding) may be affected by an application of this, or any liquid product.
3. Before use in commercial or residential food areas, refer to the Food Handling Establishment section of the label.

Remove patients in hospital, nursing homes, or any rooms before treating. If application is done as a surface spray or crack and crevice application, patients may re-enter once the spray has dried. If application is a ULV Fog or space spray, patients may return to the room after it has been thoroughly ventilated.

4. In institutions (including libraries, sports facilities, etc.) applications may be made while the facility is in use, provided treatment is not made in the immediate area where people are present.

### APPLICATION INSTRUCTIONS

PIVOT 10 may be applied both indoors and outdoors as a general surface spray, ULV/fog or space spray application, spot treatment or crack and crevice application. Application may be made using power sprayers, hand pump sprayers, void

injectors, in carpet shampoo, using a paintbrush, roller, professional foam applicator, or similar device.

**SURFACE APPLICATIONS:** Diluted spray may be applied as a spot treatment, broadcast surface application or crack and crevice spray. Apply spray solution to areas where conditions support insect populations and reproduction.

Limitations, Restrictions, and Exceptions

#### USE IN KENNELS AND DOGHOUSES

Apply as a surface application (Including Broadcast, Spot and Crack & Crevice) at the rate of 4 mL in one gallon of water (or sufficient volume of water to get adequate coverage) to treat 1,500 square feet (139 sq meter). Use up to 12 mL per 1,500 square feet (139 sq meter) for insect populations that are difficult to control or in conditions that make it difficult to get control target pests. Remove animals before spraying. Apply in and around buildings, resting areas, walls, floor, animal bedding and run areas. Animals may be returned to treated areas when spray has thoroughly dried.

Space Spray, Void Injection ULV and Fog Application: Read and Follow the \"ULV AND FOGGING APPLICATION INSTRUCTIONS, INCLUDING SPACE SPRAY AND VOID INJECTION\" To achieve uniform coverage apply PIVOT 10 (or brand name] as a ULV or fogging treatment, alone or tank mixed with an adulticide. Apply at the rate of IML of concentrate per 1,000 cu. ft. (28 cubic meter) of space to be treated. Direct application to areas where target insects are known to travel, rest, hide or harbor.

Method

[Spot treatment](#)

[Surface](#)

[Broadcast](#)

Rates

[field\\_rates 0](#)

•

Timings

[N. A.](#)