

COMMERCIAL TURF APPLICATIONS - COPRINUS SNOW MOLD

General Information

COMMERCIAL APPLICATIONS

GENERAL INFORMATION

This product is a broad-spectrum fungicide for use on a wide variety of turf and ornamental diseases. Apply this product using ground-based, hand-operated (push) or mechanical (power) equipment.

USE PRECAUTIONS

- For use only by certified applicators or those under their immediate supervision. Not for use on turf being grown for sale or other commercial use as sod.
- Do not apply this product through any type of irrigation system.
- Do not use on turf being grown for sale or other commercial use as sod.
- Do not apply this product by hand or with hand-held rotary spreader (e.g., bellygrinder)
- Do not graze animals on treated turf.
- Do not feed clippings to livestock or poultry.
- Use only properly calibrated application equipment when applying this product.
- Thoroughly clean all application equipment after each use.
- For application schedules appropriate to your area, contact your local State Extension Service for instructions.

RESISTANCE MANAGEMENT

In order to prevent the development of tolerant strains of fungi, use this product with other fungicides that use different modes of action. Do not use this product with products containing the active ingredient thiabendazole because thiabendazole has a similar chemistry and mode of action and can contribute to the development

of disease tolerance. This product can remain useful for disease control as long as the specified precautions are followed. A tolerant strain of fungi is most likely present if a treatment of this product is found to be ineffective and you should promptly contact your State Agricultural Experiment Station or State Cooperative Extension Service for disease identification and advice on other suitable fungicide or disease control strategies.

Limitations, Restrictions, and Exceptions

COMMERCIAL TURF APPLICATIONS

Use this product on all cool- and warm-season grasses such as Bentgrass, Bluegrass, Bermudagrass, Fescue, Ryegrass, St. Augustinegrass, Zoysiagrass, or their mixtures in the following sites:

- Commercial turf
- Residential turf
- Public turf (parks, athletic fields, schools, day care centers, etc.)
- Golf courses (greens, tees, fairways and aprons)

When used in accordance with the instructions in this label, this product is not phytotoxic to any of the above-mentioned grasses.

Application Instructions

- Apply under dry or moist leaf/soil conditions.
- For best results, activate the application within 24 hours with irrigation or natural rainfall. If activating by irrigation, apply 0.10 to 0.25 inch of water.
- Treat aprons when treating golf greens.

Coprinus Snow Mold: Make two treatments at 21-day intervals in late fall to early winter. Make the last application just prior to first permanent snow cover.

NOTE: Rotate with PCNB.

Refer to the label for the maximum treatment and seasonal application rates allowed based on the use site to which product is being applied.

Minimum Retreatment Interval: 14 days

Method

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[Make two treatments at 21-day intervals in late fall to early winter.](#)