

GRASS PERIMETER APPLICATION - PILL BUGS SOWBUGS AND EUROPEAN CRANE FLIES

General Information

DIRECTIONS FOR USE

Do not apply more than 0.2 lb. active ingredient (a.i.) per acre per application on residential use sites (i.e., around private homes, apartment buildings, condominiums, non-agricultural outbuildings, non-commercial greenhouses, pre-schools or day care facilities). May be applied at up to 0.4 lb. a.i. per acre per application on non-residential use sites (i.e., around institutional, public, commercial or industrial buildings; parks; recreational areas or athletic fields).

Aerial application is prohibited.

Do not broadcast apply more than 0.4 lb. a.i. (200 lbs. UP-Star GC Granular) per acre per year.

Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

Not for use on nurseries, in commercial greenhouses or on grass grown for seed.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Limitations, Restrictions, and Exceptions

GRASS PERIMETER APPLICATION

Broadcast UP-Star GC Granular Insecticide with application equipment which will ensure uniform coverage of the treatment area.

APPLICATION RATES

The application rates listed are expected to provide excellent control of the target

pests under normal conditions. However, based on the experience of the applicator, UP-Star GC Granular may be applied at up to 0.4 lb. a.i./A to control each of the pests listed. For residential lawn use, do not apply more than 0.2 lb. a.i. per acre per application.

Instructions:

In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of UP-Star GC Granular if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

Method

[Broadcast](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

[field_rates 2](#)

•

Timings

[N.A.](#)