

TREE CROPS (CITRUS, ORANGE, ETC.)

General Information

USE PRECAUTIONS

Read entire label before using. Use only as a baiting treatment to be applied directly to the soil area beneath plants as listed on the label. Do not allow bait granules to come in contact with parts of plant that may be used for food or feed purposes. The grower is responsible for residues on his crops, as well as for damages caused by drift from his property to that of others. Do not allow drift to adjoining food, fiber or pasture crops.

NOTE: TRAIL'S END LG should never be applied to dry soil.

Applications should be made following irrigation for best results. Apply any time of the day but evening applications are preferred. Do not rewater for 48 hours after application.

GENERAL INFORMATION

Snails and slugs are nocturnal and generally feed during the night damaging many varieties of plants and plant seedlings. They inhabit damp, moist areas around decaying refuse, organic matter, beneath boards, rocks and trash piles and hide at the base of growing plants.

Their presence can be detected by the shiny trails left on the soil surface, sidewalks, driveways, etc.

APPLICATION RATE

For control of SNAILS and SLUGS in the following areas, use one pound to treat 1,100 to 1,500 sq. ft. of soil surface equivalent to 30 to 40 pounds of granules per acre, unless otherwise noted below. Apply when first plant damage appears or is anticipated and repeat as needed for control.

Limitations, Restrictions, and Exceptions

TREE CROPS

Broadcast liberally under trees, around irrigation pipes, farm buildings and any other place where SLUGS and SNAILS may congregate. Citrus (Oranges, Lemons, Grapefruit, etc.): Apply up to 40 pounds per acre.

Method

[Broadcast](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[When first plant damage appears or is anticipated.](#)