

# **BANANA - ROOT DISEASE COMPLEX (POST-HARVEST PSEUDOSTEM (STUMP) INJECTION)**

## General Information

### CROP USES

Legion 80 WDG is a systemic fungicide that, when used according to the label directions and as part of good management practices programs, will control or suppress listed fungal diseases in a variety of crops.

### TOLERANCE

Although Legion 80 WDG has been found to be safe to use with the species listed on the label, it is impossible to test all species or varieties under all types of growing conditions for tolerance to Legion 80 WDG. Always test for plant phytotoxicity on a small scale using the recommended rates before preparing spray solutions or applying to large areas of plants.

### APPLICATION DIRECTIONS FOR FOLIAR SPRAYS

For systemic control of various fungal diseases, apply Legion 80 WDG according to these label directions. Apply Legion WDG in a sufficient amount of water to ensure a thorough coverage of plant foliage. The spray volumes will vary by crop and growth stage. For vegetables and small fruit, apply Legion 80 WDG by ground in spray volumes of 20 gals. or more per acre. For aerial applications to those labeled crops, apply Legion 80 WDG in spray volumes of 10 gals. or more per acre. For tree fruit, apply Legion 80 WDG until the foliage is completely wet.

### APPLICATION DIRECTIONS FOR TRUNK SPRAYS AND PAINTS

Legion 80 WDG mixed with a small volume of water may be applied as a spray or paint to wounded areas of limbs or trunks of trees (refer to the use directions for citrus and stone fruit below for rates and volumes). Use sufficient volumes of water to ensure thorough coverage of the area to be treated.

### Limitations, Restrictions, and Exceptions

## BANANA - ROOT DISEASE COMPLEX (POST-HARVEST PSEUDOSTEM (STUMP) INJECTION)

Specific Directions: To calculate the correct unit dosage required, divide the per acre rate by the number of production units per year. Use a self-refilling injector (e.g., Duomat Injector) to inject the unit dosage dissolved in water into the pseudostem (stump) that remains after the banana bunch is harvested (within one week after harvest). Make injections 3 feet above the ground.

### RESTRICTIONS:

- Do not apply in less than 5 gallons of water per acre.
- Not registered for use in California on bananas.

### Method

[Injection](#)

### Pre-Harvest Interval

1 day

### Rates

[field rates 0](#)

•

### Restricted Entry Interval

12 hours

### Timings

[Post-harvest](#)