

TOBACCO - BLUE MOLD (FOLIAR)

General Information

CROP USES

Legion 80 WDG is a systemic fungicide that, when used according to the label directions and as part of good management practices programs, will control or suppress listed fungal diseases in a variety of crops.

TOLERANCE

Although Legion 80 WDG has been found to be safe to use with the species listed on the label, it is impossible to test all species or varieties under all types of growing conditions for tolerance to Legion 80 WDG. Always test for plant phytotoxicity on a small scale using the recommended rates before preparing spray solutions or applying to large areas of plants.

APPLICATION DIRECTIONS FOR FOLIAR SPRAYS

For systemic control of various fungal diseases, apply Legion 80 WDG according to these label directions. Apply Legion WDG in a sufficient amount of water to ensure a thorough coverage of plant foliage. The spray volumes will vary by crop and growth stage. For vegetables and small fruit, apply Legion 80 WDG by ground in spray volumes of 20 gals. or more per acre. For aerial applications to those labeled crops, apply Legion 80 WDG in spray volumes of 10 gals. or more per acre. For tree fruit, apply Legion 80 WDG until the foliage is completely wet.

APPLICATION DIRECTIONS FOR TRUNK SPRAYS AND PAINTS

Legion 80 WDG mixed with a small volume of water may be applied as a spray or paint to wounded areas of limbs or trunks of trees (refer to the use directions for citrus and stone fruit below for rates and volumes). Use sufficient volumes of water to ensure thorough coverage of the area to be treated.

Limitations, Restrictions, and Exceptions

TOBACCO - BLUE MOLD (FOLIAR)

Application Method: Foliar - greenhouse, planthouse, outdoor transplant beds

Specific Directions: Make applications of Legion 80 WDG before or at first signs of Blue Mold and repeat applications every 5-7 days. Do not make more than 2 applications. Do not apply more than 0.6 lb. product per 1000 sq. ft. per application or more than 1.2 lbs. product per 1000 sq. ft. per season.

NOTE: Phytotoxicity (leaf burn) will occur in greenhouse plants if Legion 80 WDG is washed into the root zone or float water when insecticides such as the neonicotinoids (Group 4A; ex. imidacloprid) are used at the same time. THE USER ASSUMES ALL RISKS WITH USE OF THIS PRODUCT IN THE GREENHOUSE. Follow these directions for use carefully to avoid crop damage.

Use low volume sprays: For small plants, use 3 gals. of spray solution per 1000 sq. ft.; for larger plants, use 12 gals. of spray solution per 1000 sq. ft. Cover the foliage with spray, but do not allow sprays to reach the soil or float water.

Irrigate the day after application: This delay allows the Legion 80 WDG to be absorbed into the plant through the foliage sprays. If an insecticide which must reach the soil to be active must be used, apply that product before the Legion 80 WDG application.

RESTRICTIONS:

- Do not apply by air in less than 10 gallons of water per acre.
- To avoid phytotoxicity, do not use on shade tobacco.
- Not registered for use in California on tobacco.

NOTE: Tobacco may be sensitive to combinations of Legion 80 WDG with other products, especially copper, adjuvants, and foliar fertilizers or to unbuffered Legion 80 WDG application to foliage bearing copper residues. Crop sensitivity to particular combinations should be determined by spraying a small area of crop foliage and/or fruit and evaluating treated area in 3 to 7 days for adverse effects. Carefully read the sections on tank mixes, phytotoxicity, and compatibility on the label.

Method

[Foliar spray](#)

Pre-Harvest Interval

3 days

Rates

[field_rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[At first signs of Blue Mold.](#)