

# CITRUS

## General Information

### PRECAUTIONS AND RESTRICTIONS

- This product is intended solely for use on agricultural crops grown for commercial or research purposes, and in State and/or Federal invasive mollusk eradication operations. This product may not be applied to residential sites (i.e., in yards, gardens, around homes and apartments) and other sites such as the exterior areas around schools and daycare facilities, and athletic fields, playgrounds, parks, recreation areas, etc.
- Broadcast applications and applications of this product directed to plant parts are prohibited except to grass grown for seed, and as specified for use in State and/or Federal invasive mollusk eradication operations.
- Do not allow bait pellet to come in contact with plant parts.
- Use only as a baiting treatment to be applied directly to the soil area beneath listed plants.
- Read entire label before using.
- Use a band treatment between rows.
- Do not apply as a wide-area broadcast.
- Grower is responsible for residues on crops.
- Do not apply to dry soil. Make applications following irrigation.
- Do not re-water for 48 hours after application.
- Apply any time of day, but evening applications are preferred.

BLUE BOMBSHELL is a soil surface treatment for control of snails and slugs in citrus groves, berry crops, listed vegetable crops, ornamental gardens, and seed grasses. When applied as directed,

BLUE BOMBSHELL will withstand rain and sprinkling and will offer protection to gardens for several weeks. This product may be injurious to Daylilies and Clematis. Do not use on these plants.

For use on ornamentals, artichokes, blueberries, caneberries, strawberries, lettuce, cole crops and other leafy greens, and tomatoes apply in a band, between rows and/or directed towards base of the plants, by hand or with commercial application equipment designed for banded application of granular products.

For use on citrus, apply in a band under tree canopy, by hand or with commercial application equipment designed for banded application of granular products.

#### Limitations, Restrictions, and Exceptions

##### CITRUS

Minimum Days Between Retreatment: 14

##### Method

[Directed](#)

[Band](#)

##### Rates

[field rates 0](#)

- 

##### Restricted Entry Interval

12 hours

##### Timings

[Apply any time of day, but evening applications are preferred.](#)