

TREES AND SHRUBS: NON-BEARING CITRUS

General Information

DIRECTIONS FOR USE

Not for use on plants being grown for sale or for other commercial use, or for commercial seed production or for research purposes.

Step #1: For vegetables, measure the area to be sprayed. Calculate square feet by multiplying length by width. For all other applications, apply as a thorough cover spray.

Step #2: Use a clean sprayer. Carefully measure and mix the amount of concentrate and water as directed.

Step #3: Spray as directed. Thorough coverage is important. Flush sprayer with clean water after each use.

Make applications when insects first appear. Repeat applications as directed. See directions for complete insect list.

People and pets should not be allowed in treated areas until spray has dried.

GENERAL DIRECTIONS

Use 2 fl. oz. in 1 gal. of water. Apply when insects appear. Wet the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage.

Limitations, Restrictions, and Exceptions

TREES & SHRUBS

How to use:

Use 1 fl. oz. in 1 gal. of water. Apply as a thorough spray, wetting leaves and branches to the dripping point. Try to penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use intervals of 4-8 days. Do not exceed 16 applications per season. Application can be

made up to the day of collection.

Method

[Spray](#)

Rates

[field rates 0](#)

•

Timings

[Spray in the late afternoon or evening, when the temperature ranges from 50° to 75°F and when there is little or no wind.](#)