

# **FOLIAR APPLICATION (VEGETABLES AND POTATOES)**

## General Information

### PRODUCT INFORMATION

BIOMIN MAGNESIUM contains nitrogen and chelated magnesium for soil and/or foliar applications. It is completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. BIOMIN MAGNESIUM is designed to prevent and correct magnesium deficiency and boost crops during critical or fast growing periods.

BIOMIN MAGNESIUM may be applied with nitrogen fertilizer.

BIOMIN MAGNESIUM can be applied to ALL commercial crops including all vegetables, Potatoes, Carrots, Onions, all berries and strawberries, all Brassicas, all fruit trees, Avocados, Persimmons, Citrus, all Nut trees, Vineyards, Kiwifruit, Tropical fruit, Lettuce, Asparagus, all Melons, Beans, Ginger, Ornamentals, Cut flowers and Pastures.

BIOMIN MAGNESIUM is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 200 times of water. For best results and more specific recommendations, contact your nearest distributor. Always perform a compatibility test before mixing any products with BIOMIN MAGNESIUM.

## Limitations, Restrictions, and Exceptions

### FOLIAR APPLICATION

Vegetables and Potatoes: Apply 1.0-1.5 pounds of BIOMIN MAGNESIUM per acre (1.1-1.7 Kg/Ha) at about 2-4 weeks after emergence or if deficiency symptoms appear. Repeat the application 2-3 weeks later.

#### Method

[Foliar spray](#)

#### Rates

[field rates 0](#)

[field rates 1](#)

-

## Timings

At about 2-4 weeks after emergence or if deficiency symptoms appear.