

## **FOLIAR APPLICATION (GREENHOUSE CROPS)**

### General Information

#### PRODUCT INFORMATION

BIOMIN MAGNESIUM contains nitrogen and chelated magnesium for soil and/or foliar applications. It is completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. BIOMIN MAGNESIUM is designed to prevent and correct magnesium deficiency and boost crops during critical or fast growing periods.

BIOMIN MAGNESIUM may be applied with nitrogen fertilizer.

BIOMIN MAGNESIUM can be applied to ALL commercial crops including all vegetables, Potatoes, Carrots, Onions, all berries and strawberries, all Brassicas, all fruit trees, Avocados, Persimmons, Citrus, all Nut trees, Vineyards, Kiwifruit, Tropical fruit, Lettuce, Asparagus, all Melons, Beans, Ginger, Ornamentals, Cut flowers and Pastures.

BIOMIN MAGNESIUM is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 200 times of water. For best results and more specific recommendations, contact your nearest distributor. Always perform a compatibility test before mixing any products with BIOMIN MAGNESIUM.

### Limitations, Restrictions, and Exceptions

#### FOLIAR APPLICATION

Greenhouse Crops: Apply 1.0-1.5 pounds of BIOMIN MAGNESIUM per acre (1.1-1.7 Kg/Ha) 2-3 weeks after emergence or transplanting, or when boron deficiencies first appear. Repeat the application every one or two weeks.

#### Method

[Foliar spray](#)

#### Rates

[field rates 0](#)

[field rates 1](#)

-

Timings

2-3 weeks after emergence or transplanting, or when magnesium deficiencies first appear.