

## **LEAFY VEGETABLES (EXCEPT BRASSICA VEGETABLES)**

### General Information

### DIRECTIONS FOR USE

#### Ground Application

Apply the specified rate of this product in sufficient water for thorough coverage of foliage for optimum disease control.

#### Aerial Application

Apply the specified rate of this product in 3–5 gallons of water per acre for field crops. For tree crops, apply in no less than 10 gallons per acre.

### USE INFORMATION

Phiticide is a systemic product containing mono and dibasic phosphites. The phosphonic ion is effective in controlling downy mildew, brown rot, foot rot and other diseases caused by *Phytophthora*, *Pythium* and other related fungi by activating the plant's natural resistance mechanisms. When used in a complete Integrated Pest Management (IPM) disease control program with good cultural practices, this product will provide control of the listed diseases. To achieve the best results in disease control spraying, initiate before signs or symptoms of the disease are present or when the disease first appears or environmental conditions are conducive to disease development. The preharvest interval for this product is 0 days.

### Limitations, Restrictions, and Exceptions

#### LEAFY VEGETABLES (EXCEPT BRASSICA VEGETABLES)

*Bremia lactucae*, *Peronospora* spp. (Downy Mildew): Apply 2.5 to 5 pints per acre in 10 to 50 gallons of water to obtain good coverage. Begin application when conditions favor disease development and continue on 7 to 21 day spray intervals. Use higher rate and shorter spray interval during times of moderate to high disease pressure.

NOTE: Do not exceed seven applications per season.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Restricted Entry Interval

4 hours

Timings

[Begin application when conditions favor disease development.](#)