

ONIONS (DRY BULB) AND OTHER ALLIUM SPP

General Information

DIRECTIONS FOR USE

Ground Application

Apply the specified rate of this product in sufficient water for thorough coverage of foliage for optimum disease control.

Aerial Application

Apply the specified rate of this product in 3–5 gallons of water per acre for field crops. For tree crops, apply in no less than 10 gallons per acre.

USE INFORMATION

Phiticide is a systemic product containing mono and dibasic phosphites. The phosphonic ion is effective in controlling downy mildew, brown rot, foot rot and other diseases caused by *Phytophthora*, *Pythium* and other related fungi by activating the plant's natural resistance mechanisms. When used in a complete Integrated Pest Management (IPM) disease control program with good cultural practices, this product will provide control of the listed diseases. To achieve the best results in disease control spraying, initiate before signs or symptoms of the disease are present or when the disease first appears or environmental conditions are conducive to disease development. The preharvest interval for this product is 0 days.

Limitations, Restrictions, and Exceptions

ONIONS (DRY BULB) AND OTHER ALLIUM SPP.

Peronospora destructor (Downy Mildew), Alternaria porri (Purple Blotch): Best results are obtained when this product is used in a preventative disease control program. Apply 2.5 to 3.75 pints in 20 to 50 gallons of water per acre beginning when conditions first favor disease development. Continue application on 7 to 14 day intervals. If disease is already present, use the high rate and shortest spray interval.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Restricted Entry Interval

4 hours

Timings

[Beginning when conditions first favor disease development.](#)