

STRAWBERRY - FOLIAR APPLICATION (ANNUAL PLANTING)

General Information

DIRECTIONS FOR USE

Ground Application

Apply the specified rate of this product in sufficient water for thorough coverage of foliage for optimum disease control.

Aerial Application

Apply the specified rate of this product in 3–5 gallons of water per acre for field crops. For tree crops, apply in no less than 10 gallons per acre.

USE INFORMATION

Phiticide is a systemic product containing mono and dibasic phosphites. The phosphonic ion is effective in controlling downy mildew, brown rot, foot rot and other diseases caused by Phytophthora, Pythium and other related fungi by activating the plant's natural resistance mechanisms. When used in a complete Integrated Pest Management (IPM) disease control program with good cultural practices, this product will provide control of the listed diseases. To achieve the best results in disease control spraying, initiate before signs or symptoms of the disease are present or when the disease first appears or environmental conditions are conducive to disease development. The preharvest interval for this product is 0 days.

Limitations, Restrictions, and Exceptions

STRAWBERRY

Foliar Application: Annual planting — Apply 2.5 to 5.0 pints per acre. Begin application 2 to 3 weeks after planting and repeat on 30 to 60 day intervals while conditions favor disease development.

If using Red Stele susceptible varieties or if disease pressure is severe, use higher rates (3.75 to 5 pints per acre), shortest application time (30 days) and maximum number of applications.

Method

[Foliar application](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

4 hours

Timings

[2 to 3 weeks after planting.](#)