

# **TOMATO, TOMATILLO**

## General Information

### DIRECTIONS FOR USE

#### Ground Application

Apply the specified rate of this product in sufficient water for thorough coverage of foliage for optimum disease control.

#### Aerial Application

Apply the specified rate of this product in 3–5 gallons of water per acre for field crops. For tree crops, apply in no less than 10 gallons per acre.

### USE INFORMATION

Phiticide is a systemic product containing mono and dibasic phosphites. The phosphonic ion is effective in controlling downy mildew, brown rot, foot rot and other diseases caused by *Phytophthora*, *Pythium* and other related fungi by activating the plant's natural resistance mechanisms. When used in a complete Integrated Pest Management (IPM) disease control program with good cultural practices, this product will provide control of the listed diseases. To achieve the best results in disease control spraying, initiate before signs or symptoms of the disease are present or when the disease first appears or environmental conditions are conducive to disease development. The preharvest interval for this product is 0 days.

#### Limitations, Restrictions, and Exceptions

### TOMATO, TOMATILLO

Phytophthora spp. (Root Rot): Apply 2.5 to 5.0 pints in a minimum of 10 gallons of water per acre beginning at the 2 to 4 leaf growth stage for direct seeded tomatoes or immediately after transplanting to the field. Continue on 7 to 14 day spray intervals when conditions are favorable for disease development. Use the higher rate (3.75 pints per acre) and shorter spray interval when disease pressure is moderate to high.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field\\_rates 0](#)

- 

Restricted Entry Interval

4 hours

Timings

[Beginning at the 2 to 4 leaf growth stage for direct seeded tomatoes or immediately after transplanting to the field.](#)