

BEDDING PLANTS

General Information

DIRECTIONS FOR USE

Ground Application

Apply the specified rate of this product in sufficient water for thorough coverage of foliage for optimum disease control.

Aerial Application

Apply the specified rate of this product in 3–5 gallons of water per acre for field crops. For tree crops, apply in no less than 10 gallons per acre.

USE INFORMATION

Phiticide is a systemic product containing mono and dibasic phosphites. The phosphonic ion is effective in controlling downy mildew, brown rot, foot rot and other diseases caused by Phytophthora, Pythium and other related fungi by activating the plant's natural resistance mechanisms. When used in a complete Integrated Pest Management (IPM) disease control program with good cultural practices, this product will provide control of the listed diseases. To achieve the best results in disease control spraying, initiate before signs or symptoms of the disease are present or when the disease first appears or environmental conditions are conducive to disease development. The preharvest interval for this product is 0 days.

Limitations, Restrictions, and Exceptions

- Begin applications prior to disease development and in conjunction with good cultural management practices. Use the highest rate when disease pressure is severe. Do not exceed specified rates or apply more frequently than at specified intervals or plant injury may occur.

BEDDING PLANTS

FOLIAR APPLICATIONS to plants such as Begonia, Pansy, Marigold, Zinnia, Petunia, Geranium and Impatiens: Mix 1.25 to 4 pints with 100 gallons of water and spray to

wet using no more than 400 gallons of water per acre. Repeat as necessary but do not exceed one application every 14 days.

Method

[Foliar application](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

4 hours

Timings

[N.A.](#)