

TURF - YELLOW TUFT (21 DAYS APPLICATION INTERVAL)

General Information

DIRECTIONS FOR USE

Ground Application

Apply the specified rate of this product in sufficient water for thorough coverage of foliage for optimum disease control.

Aerial Application

Apply the specified rate of this product in 3–5 gallons of water per acre for field crops. For tree crops, apply in no less than 10 gallons per acre.

USE INFORMATION

Phiticide is a systemic product containing mono and dibasic phosphites. The phosphonic ion is effective in controlling downy mildew, brown rot, foot rot and other diseases caused by Phytophthora, Pythium and other related fungi by activating the plant's natural resistance mechanisms. When used in a complete Integrated Pest Management (IPM) disease control program with good cultural practices, this product will provide control of the listed diseases. To achieve the best results in disease control spraying, initiate before signs or symptoms of the disease are present or when the disease first appears or environmental conditions are conducive to disease development. The preharvest interval for this product is 0 days.

Limitations, Restrictions, and Exceptions

- Begin applications prior to disease development and in conjunction with good cultural management practices. Use the highest rate when disease pressure is severe. Do not exceed specified rates or apply more frequently than at specified intervals or plant injury may occur.

TURF

This product is a systemic fungicide, which can be used, in a seasonal program, for

the suppression of Phytophthora and Pythium diseases, such as blight and root rot, and yellow tuft on common turf grasses. Apply as a foliar spray on golf courses, sod farms and other turf areas using 1 to 5 gallons of water per 1000 square feet with a properly calibrated sprayer as indicated.

- Refer in the label for TURF TANK MIXTURES information.

Method

[Foliar spray](#)

Rates

[field_rates 0](#)

•

Restricted Entry Interval

4 hours

Timings

[N.A.](#)