

FIRE BLIGHT ON ORNAMENTALS

General Information

DIRECTIONS FOR USE

Ground Application

Apply the specified rate of this product in sufficient water for thorough coverage of foliage for optimum disease control.

Aerial Application

Apply the specified rate of this product in 3–5 gallons of water per acre for field crops. For tree crops, apply in no less than 10 gallons per acre.

USE INFORMATION

Phiticide is a systemic product containing mono and dibasic phosphites. The phosphonic ion is effective in controlling downy mildew, brown rot, foot rot and other diseases caused by Phytophthora, Pythium and other related fungi by activating the plant's natural resistance mechanisms. When used in a complete Integrated Pest Management (IPM) disease control program with good cultural practices, this product will provide control of the listed diseases. To achieve the best results in disease control spraying, initiate before signs or symptoms of the disease are present or when the disease first appears or environmental conditions are conducive to disease development. The preharvest interval for this product is 0 days.

Limitations, Restrictions, and Exceptions

FIRE BLIGHT ON ORNAMENTALS

Apply 1.25 quarts per 100 gallons of water, as a foliar spray, to thoroughly wet the foliage. Begin applications at the pre-bloom stage and continue until the end of bloom period using a 7-day intervals.

Method

[Foliar spray](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

4 hours

Timings

[At the pre-bloom stage.](#)