

# **WEEDY BRUSH, TREES AND VINES CONTROLLED**

## General Information

### GENERAL INFORMATION

#### ARBORSYSTEMS DIRECT-INJECT TREE INJECTION SYSTEMS

The product is recommended for woody brush, tree and vine control in all non-cropland sites including forestry, roadsides, rights-of-way, industrial residential, fencerows, railroads and aquatic sites. Woody vegetation may be controlled by injection of EZ-Ject Diamondback Herbicide shells.

The product moves throughout the tree from the point of injection to all roots and shoots. Symptoms appear gradually (normally within 2 to 4 weeks) beginning with wilting, yellowing and browning of the foliage. This is followed by deterioration of all roots and shoots.

For aquatic sites, the product may be injected into stems of trees and brush standing in water or wetlands. Do not apply directly to water or inject into stems below the water level.

### APPLICATION INSTRUCTIONS

Inject the shells using only EZ-Ject shell injection equipment. The shells must penetrate through the outer bark into living phloem tissue to provide effective results.

For trees with less than 2.5 inch DBH (diameter at breast height), inject one shell per stem. Inject one shell every 4 inches, evenly around the trunk of the tree, below all major branches. Tree may be injected at any time of the year except when wood is frozen.

Method

[Injection](#)

Timings

[N.A.](#)