

NON-CROPLAND - ANNUAL/PERENNIAL CONTROL

General Information

PRECAUTIONS AND RESTRICTIONS

Do not apply the product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Aerial application is prohibited.

For turf use, the maximum number of broadcast applications is two per year.

Do not use the product in or near greenhouses.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product runoff, do not overwater the treated area(s) or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

Avoid broadcast and small spot treatment applications when temperatures exceed 90°F, as turf injury may occur.

Avoid applying during excessively dry or hot periods unless irrigation is used. For optimum results, turf should not be mowed for 1 to 2 days before and after application, do not apply if rain is expected within 4 hours after the application, and delay irrigation cycle for 24 hours. Reseed no sooner than 3 to 4 weeks after application of the product. Failure to observe all precautions may result in injury to turf and/or susceptible plants.

Avoid mist to vegetables, flowers, ornamentals, shrubs, trees, and other desirable plants. Do not pour spray solutions near these plants.

Susceptible Plants: Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans; and other

vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements: Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment: All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional requirements for ground boom application:

Do not apply with a nozzle height greater than 4 feet above the application site.

Adjuvant Use: When an adjuvant is to be used with the product, Control Solutions, Inc. suggests the use of a Chemical Producers and Distributors Association certified adjuvant.

Drift Control And Defoaming: For drift control and defoaming, use a quality defoamer.

Fertilizer Use: The product can be mixed with some liquid fertilizers or liquid iron metals. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration, and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All state and/or federal regulations relating to the application of liquid fertilizers or liquid iron and the product must be strictly followed.

The product will kill, control, or suppress the weeds listed in the label for the product. Some of these species may require repeat spot applications even under ideal conditions for application.

For use in New York State by spot treatment only. Spray only the weed not the entire lawn.

Limitations, Restrictions, and Exceptions

NON-CROPLAND (fence rows, hedge rows, roadsides, ditches, rights-of-way, utility, power lines, railroads, airports, and industrial sites)

Applications to non-cropland areas are not applicable to treatment of commercial

timber or other plants being grown for sale or other commercial use, for commercial seed production, or for research purposes.

Roadsides (including aprons and guardrails), rights-of-way and other similar noncrop areas: For control of broadleaf weeds, mix at a rate of 1 to 2 quarts of the product per acre in enough water to wet all parts of the foliage. This may require 50 to 300 gallons of water per acre. This mixture will cover 43,560 square feet. Thoroughly saturate all weeds with spray mixture. Apply anytime between the time when the plants come into full leaf (spring) to when the plants begin to go dormant. Best results are obtained when the weeds are young and actively growing. Do not cut weeds until herbicide is translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with the high water volume. For small (spot) applications with small tank sprayers, apply at the rate of 4 ounces of the product per gallon of water and spray to thoroughly wet all foliage.

RESTRICTIONS FOR NON-CROPLAND

Annual/Perennial Control (Postemergence):

- Minimum of 30 days between applications.

Method

[Broadcast/Foliar Ground](#)

[Spot treatment](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

48 hours

Timings

[Postemergence \(Weed\)](#)